GUIDE
FOR THE CARE AND USE OF
LABORATORY ANIMALS

Eighth Edition

THE NATIONAL ACADEMIES PRESS 500 Fifth Street, NW Washington, DC 20001

NOTICE: The project that is the subject of this report was approved by the Governing Board of the National Research Council, whose members are drawn from the

The National Academy of Sciencesis a private, nonpro t, self-perpetuating society of distinguished scholars engaged in scienti c and engineering research, dedicated to the furtherance of science and technology and to their use for the general welfare. Upon the authority of the charter granted to it by the Congress in 1863, the Acad

COMMITTEE FOR THE UPDATE OF THE GUIDE FOR THE CARE AND USE OF LABORATORY ANIMALS

Members

Janet C. Garber (Chair), Garber Consulting
R. Wayne Barbee, Virginia Commonwealth University
Joseph T. Bielitzki, University of Central Florida
Leigh Ann Clayton, National Aquarium, Baltimore
John C. Donovan, BioResources, Inc.
Coenraad F. M. Hendriksen, Netherlands Vaccine Institute, Bilthoven, The Netherlands (until March 2009)
Dennis F. Kohn, Columbia University (retired)
Neil S. Lipman, Memorial Sloan-Kettering Cancer Center and Weill Cornell Medical College
Paul A. Locke, Johns Hopkins Bloomberg School of Public Health
John Melcher, U.S. Senate (retired)
Fred W. Quimby, Rockefeller University (retired)
Patricia V. Turner, University of Guelph, Canada
Geoffrey A. Wood, University of Guelph, Canada
Hanno Würbel, Justus Liebig University of Giessen, Germany

Staff

Lida Anestidou, Study Director
Frances Sharples, Acting Director
Kathleen Beil, Administrative Coordinator
Cameron H. Fletcher, Senior Editor
Ruth Crossgrove, Senior Editor
Radiah Rose, Manager of Editorial Projects
Rhonda Haycraft, Senior Project Assistant
Joanne Zurlo, Director (until April 2010)

INSTITUTE FOR LABORATORY ANIMAL RESEARCH COUNCIL

Members

Stephen W. Barthold (Chair), Center for Comparative Medicine, University of California-Davis
Kathryn A. Bayne, Association for Assessment and Accreditation of Laboratory Animal Care International, Frederick, Maryland
Myrtle A. Davis, National Cancer Institute, National Institutes of Health, Bethesda, Maryland
Jeffrey I. Everitt, Comparative Medicine and Investigator Support, GlaxoSmithKline Research and Development, Research Triangle Park, North Carolina (until June 2010)
James G. Fox, Division of Comparative Medicine, Massachusetts Institute of Technology, Cambridge
Nelson L. Garnett, Laboratory Animal Care and Use Programs, Dickerson, MD
Estelle B. Gauda, Johns Hopkins University School of Medicine, Johns Hopkins Hospital, Baltimore, Maryland (until June 2010)
Joseph W. Kemnitz, Institute for Clinical and Translational Research and Department of Physiology, University of Wisconsin-Madison
Judy A. MacArthur Clark, Animals in Scientific Procedures Inspectorate, Home Office, London, United Kingdom
Martha K. McClintock, Institute for Mind and Biology, University of Chicago, Illinois
Leticia V. Medina, Animal Welfare and Compliance, Abbott Laboratories, Abbott Park, Illinois
Timo Olavi Nevalainen

Staff

Frances Sharples, Acting Director
Lida Anestidou, Senior Program Officer
Kathleen Beil, Administrative Coordinator
Cameron H. Fletcher, Managing Editor, ILAR Journal
Rhonda Haycraft, Program Associate
Joanne Zurlo, Director (until April 2010)

Reviewers

This eighth edition of the Guide for the Care and Use of Laboratory Animals has been reviewed in draft form by individuals chosen for

Gemma Perretta, National Research Council, Italy
Marky E. Pitts, IACUC Consultant
George E. Sanders, University of Washington
Allen W. Singer, Battelle Memorial Institute

included research scientists, veterinarians, and nonscientists representing biomedical ethics and the public's interest in animal welfare. The Committee widely solicited written and oral comments on the update of the Guide from the scientific community and the general public; comments at open meetings (on September 26, 2008, in Washington, DC; October 16, 2008, in Irvine, California; and November 14, 2008, in Chicago) as well as written comments submitted to or requested by the Committee were considered. In addition, the Committee studied the materials submitted to NIH in response to its 2005 Request for Information (NOT-OD-06-011). All comments contributed substantially to this eighth edition of the Guide.

In approaching its task, the Committee carried forward the balance between ethical and science-based practice that has always been the basis of the Guide, and fulfilled its role to provide an updated resource that enables the research community to proceed responsibly and in a self-regulatory manner with animal experimentation. The Guide is predicated on the understanding that the exercise of professional judgment both upholds the central notion of performance standards and obviates the need for more stringent regulations.

Laboratory animal science is a rapidly evolving field and the Committee identified a number of areas in which current available scientific information is insufficient; additional objective information and assessment are needed to provide a scientific basis for recommendations in future editions of the Guide. Although pursuing these concepts was beyond this Committee's charge, the following two topics merit further study: (1) space and housing needs of laboratory species and (2) the need and best methods for providing enrichment, exercise, and human contact.

The need for continual updating of the Guide is implicit in its objective "to provide information that will enhance animal well-being, the quality of research, and the advancement of scientific knowledge that is relevant to both humans and animals" (Chapter 1). The irregular and increasing intervals between updates, reaching a 14-year gap between the seventh edition and this eighth edition, mean that important new research findings might wait more than a decade before being reflected in recommended practice. Addressing this concern was beyond the charge of this Committee; we noted, however, that regular and more frequent updates of the information in the Guide will promote laboratory animal welfare and support high-quality scientific data. A formal process for revising the information in the Guide, including the updating of practice standards, could meet this need.

In undertaking this update, the Committee acknowledged the contributions of William I. Gay and Bennett J. Cohen in the development of the original Guide. In 1959, Animal Care Panel (ACP) President Cohen appointed the Committee on Ethical Considerations in the Care of Laboratory Animals to evaluate animal care and use. That Committee was chaired by Dr. Gay,

who soon recognized that the Committee could not evaluate animal care programs objectively without appropriate criteria on which to base its evaluations—that is, standards were needed. The ACP Executive Committee agreed, and the Professional Standards Committee was appointed. NIH later awarded the ACP a contract to "determine and establish a professional standard for laboratory animal care and facilities." Dr. Cohen chaired the ACP Animal Facilities Standards Committee, which prepared the first Guide for Laboratory Animal Facilities and Care.

This edition of the Guide was financially supported by the National Institutes of Health; the Office of Research Integrity, Department of Health and Human Services; the US Department of Agriculture (USDA); the Association for Assessment and Accreditation of Laboratory Animal Care International; the American Association for Laboratory Animal Science; Abbott Fund; Pfizer, Inc.; the American College of Laboratory Animal Medicine; the American Society of Laboratory Animal Practitioners; and the Association of Primate Veterinarians.

The Committee for the Update of the Guide for the Care and Use of Laboratory Animals

Overview

This eighth edition of the Guide is divided into five chapters and four appendices.

Chapter 1 presents the goals and intended audiences of the Guide as well as key concepts and terminology essential to its premise and use. Incorporating some of the material from the Introduction to the last edition, the chapter highlights a commitment to the concepts of the Three Rs—Replacement, Reduction, and Refinement—and provides an enhanced discussion of the ethics of animal use and investigator/institutional obligations.

Chapter 2 focuses on the overall institutional animal care and use program. Chapter 2 a(s)-3(o)-3(m)-3(e)13(g)-3(ram)-3(e)13(n)-3(n)-3(w-)-3(e)13((n)-3(d)

ing the growing role of aquatic animals in biomedical research. The chapter provides recommendations for housing and environment, discusses the importance of social housing, and includes enhanced sections on environmental enrichment, animal well-being, and scientific validity.

Space recommendations were nominally expanded based on the Committee's professional and expert opinion and on current housing methods. Cage sizes have historically been interpreted as minimum space needs by users of the *Guide*, and were labeled as such ("recommended minimum space") in this edition. The use of the word "minimum" does not further restrict users of the *Guide* because, although the space requirements are numbers (i.e., engineering standards), they are used in a performance standards framework. The Committee recommends minimum space for female rodents with litter and an increase of the cage height for rabbits to 16". Further, in light of many comments submitted to the Committee requesting more information on performance goals and how to achieve them, rodent breeding recommendations are accompanied by substantial guidance.

With respect to nonhuman primates (NHPs), the Committee endorses social housing as the default and has provided some species-specific guidance. An additional group has been added for monkeys, and chimpanzees

Contents

Personnel Security, 23
Investigating and Reporting Animal Welfare Concerns, 23
Program Oversight, 24
The Role of the IACUC, 24
IACUC Constitution and Function, 24
Protocol Review, 25
Special Considerations for IACUC Review, 27

Aquatic Management, 84
Behavior and Social Management, 84
Husbandry, 84
Population Management, 87
References, 88

4 VETERINARY CARE 105
Animal Procurement and Transportation, 106
Animal Procurement, 106
Transportation of Animals, 107
Preventive Medicine, 109
Animal Biosecurity, 109
Quarantine and Stabilization, 110
Separation by Health Status and Species, 111
Surveillance, Diagnosis, Treatment, and Control of Disease, 112
Clinical Care and Management, 113
Medical Management, 114
Emergency Care, 114
Recordkeeping, 115
Surgery, 115
Training, 115
Presurgical Planning, 116
Surgical Facilities, 116
Surgical Procedures, 117
Aseptic Technique, 118
Intraoperative Monitoring, 119
Postoperative Care, 119
Pain and Distress, 120
Anesthesia and Analgesia, 121
Euthanasia, 123
References, 124

5 PHYSICAL PLANT 133
General Considerations, 133
Location, 134
Centralization Versus Decentralization, 134
Functional Areas, 135
Construction Guidelines, 136
Corridors, 136
Animal Room Doors, 137
Exterior Windows, 137
Floors, 137

Drainage, 138
Walls and Ceilings, 138
Heating, Ventilation, and Air Conditioning (HVAC), 139
Power and Lighting, 141
Storage Areas, 141
Noise Control, 142
Vibration Control, 142
Facilities for Sanitizing Materials, 143
Environmental Monitoring, 143
Special Facilities, 144
Surgery, 144
Barrier Facilities, 145
Imaging, 146
Whole Body Irradiation, 147
Hazardous Agent Containment, 148
Behavioral Studies, 149
Aquatic Species Housing, 150
Security and Access Control, 151
References, 151

ADDENDUM 155

APPENDICES

A ADDITIONAL SELECTED REFERENCES 161
Subject Matter, 161
Use of Laboratory Animals, 162
Alternatives, 162
Ethics and Welfare, 163
Experimental Design and Statistics, 164
Research and Testing Methodology, 165
Program Management, 167
General References, 167
Laws, Regulations, and Policies, 168
Education, 169
Monitoring the Care and Use of Animals, 169
Occupational Health and Safety, 170
Environment, Housing, and Management, 172
General References, 172
Environmental Enrichment, 173
Genetics and Genetically Modi ed Animals, 175

Species-Speci c References—Environment, Housing, and Management, 176
Agricultural Animals, 176
Amphibians, Reptiles, and Fish, 178
Birds, 179
Cats and Dogs, 180
Exotic, Wild, and Zoo Animals, 181
Nonhuman Primates, 182
Rodents and Rabbits, 184
Other Animals, 187
Veterinary Care, 188
Transportation, 188
Anesthesia, Pain, and Surgery, 188
Disease Surveillance, Diagnosis, and Treatment, 190
Pathology, Clinical Pathology, and Parasitology, 190
Species-Speci c References—Veterinary Care, 191
Agricultural Animals, 191
Amphibians, Reptiles, and Fish, 192
Birds, 193
Cats and Dogs, 193
Exotic, Wild, and Zoo Animals, 193
Nonhuman Primates, 194
Rodents and Rabbits, 194
Design and Construction of Animal Facilities, 196

B U.S. GOVERNMENT PRINCIPLES FOR THE UTILIZATION AND CARE OF VERTEBRATE ANIMALS USED IN TESTING, RESEARCH, AND TRAINING 199

C STATEMENT OF TASK 201

D ABOUT THE AUTHORS 203

INDEX 209

1

Key Concepts

This edition of the Guide for the Care and Use of Laboratory Animals (the Guide) strongly af rms the principle that all who care for, use, or produce animals for research, testing, or teaching must assume responsibility for their well-being. The Guide is created by scientists and veterinarians for scientists and veterinarians to uphold the scienti c rigor and integrity of biomedical research with laboratory animals as expected by their colleagues and society at large.

The Guide plays an important role in decision making regarding the use of vertebrate laboratory animals because it establishes the minimum ethical, practice, and care standards for researchersarding g ginT-3(r)15(c)-3(h)-3(e)

Taken e,ether, ee practical effect of eese laws, egul(c)-3(e)-3ir a and pol cies s e,stablish ystem f elf-regul(c)-3(e)-3ir nd regul(c)-3(e)-3(,)-3( )8<us(mversight bus(mpractical,s(methical,s(mand s-3(sc)-3(i)0jee-323nti cs(mprinciles.(r9(T)-3(h)-3(i) establishes ig(,)-3( )8<ous rogram f nsmal are nd se nd rovsdes ex-]TJ EMC /Spa

scope and nature of this responsibility can vary based on the scienti c discipline, nature of the animal use, and species involved, but because it affects animal care and use in every situation this responsibility requires that producers, teachers, researchers, and institutions carry out purposeful analyses of proposed uses of laboratory animals. The Guide is central to these analyses and to the development of a program in which humane care is incorporated into all aspects of laboratory animal care and use.

APPLICABILITY AND GOALS

In the Guide, laboratory animals (also referred to as animals) are gener- ally de ned as any vertebrate animal (i.e., traditional laboratory animals, agricultural animals, wildlife, and aquatic species) produced for or used in research, testing, or teaching. Animal use is de ned as the proper care, use, and humane treatment of laboratory animals produced for or used in research, testing, or teaching.



important to keep in mind that the Guide is intended to provide information to assist researchers, institutional animal care and use committees (IACUCs), veterinarians, and other stakeholders in ensuring the implementation of effective and appropriate animal care and use programs that are based on humane care. Throughout the Guide, scientists and institutions are encouraged to give careful and deliberate thought to the decision to use animals, taking into consideration the contribution that such use will make to new knowledge, ethical concerns, and the availability of alternatives to animal use (NRC 1992). A practical strategy for decision making, the "Three Rs" (Replacement, Reduction, and Re nement) approach, is discussed in more detail below. Institutions should use the recommendations in the Guide as a foundation for the development of a comprehensive animal care and use program and a process for continually improving this program.

INTENDED AUDIENCES AND USES OF THE GUIDE

The Guide is intended for a wide and diverse audience, including

- the scienti c community
- administrators
- IACUCs
- veterinarians
- educators and trainers
- producers of laboratory animals
- accreditation bodies
- regulators
- the public.

The Guide is meant to be read by the user in its entirety, as there are

valuable for their oversight and support of animal care and use. Educators and trainers can use the Guide as a document to assess both the scope and adequacy of training programs supported by the institution. Accreditation bodies will find the Guide useful for evaluating many areas of animal care and use programs not subject to strict engineering standards (see definition below). Finally, members of the public should feel assured that adherence to the Guide will ensure humane care and use of laboratory animals.

Readers are reminded that the Guide is used by a diverse group of national and international institutions and organizations, many of which are covered by neither the Animal Welfare Act nor the PHS Policy. The Guide uses some terminology that is both defined by US statute and denotes a general concept (e.g., "attending veterinarian," "adequate veterinary care," and "institutional official"). Even if these terms are not consistent with those used by non-US institutions, the underlying principles can still be applied. In all instances where Guide recommendations are different from applicable legal or policy requirements, the higher standard should apply.

ETHICS AND ANIMAL USE

The decision to use animals in research requires critical thought, judgment, and analysis. Using animals in research is a privilege granted by society to the research community with the expectation that such use will provide either significant new knowledge or lead to improvement in human and/or animal well-being (McCarthy 1999; Perry 2007). It is a trust that mandates responsible and humane care and use of these animals. The Guide endorses the responsibilities of investigators as stated in the U.S. Government Principles for Utilization and Care of Vertebrate Animals Used in Testing, Research, and Training (IRAC 1985; see Appendix B). These principles direct the research community to accept responsibility for the care and use of animals during all phases of the research effort. Other government agencies and professional organizations have published similar principles (NASA 2008; NCB 2005; NIH 2006, 2007; for additional references see Appendix A). Ethical considerations discussed here and in other sections of the Guide should serve as a starting

design in laboratory animal research (Russell and Burch 1959). Over the years, the Three Rs have become an internationally accepted approach for researchers to apply when deciding to use animals in research and in designing humane animal research studies.

Replacement refers to methods that avoid using animals. The term includes absolute replacements (i.e., replacing animals with inanimate systems such as computer programs) as well as relative replacements (i.e., replacing animals such as vertebrates with animals that are lower on the phylogenetic scale).

Re nement refers to modi cations of husbandry or experimental prored(p)-3(e)-(

Humane Care

Humane care means those actions taken to ensure that laboratory animals are treated according to high ethical and scienti c standards. Implementation of a humane care program, and creation of a laboratory environment in which humane care and respect for animals are valued and encouraged, underlies the core requirements of the Guide and the system

expertise, and research goals) so that implementation can be best tailored to meet the recommendations in the *Guide*.

Ideally, engineering and performance standards are balanced, setting a target for optimal practices, management, and operations while encouraging flexibility and judgment, if appropriate, based on individual situations (Gonder et al. 2001).

Scientists, veterinarians, technicians, and others have extensive experience and information covering many of the topics discussed in the *Guide*. For topics on which information is insufficient or incomplete, sustained research into improved methods of laboratory animal management, care, and use is needed for the continued evaluation and improvement of performance and engineering standards.

Practice standard means the application of professional judgment by qualified, experienced individuals to a task or process over time, an approach that has been demonstrated to benefit or enhance animal care and use. Professional judgment comes from information in the peer-reviewed scientific literature and textbooks and, as in many other disciplines, from time-proven experiences in the field (for additional information see Chapter 2). In the absence of published scientific literature or other definitive sources, where experience has demonstrated that a particular practice improves animal care and use, practice standards have been used in determining appropriate recommendations in the *Guide*. In most situations, the *Guide* is intended to provide flexibility so that institutions can modify practices and procedures with changing conditions and new information.

POLICIES, PRINCIPLES, AND PROCEDURES

Policies commonly derive from a public agency or private entity. They are generally practical statements of collective wisdom, convention, or management direction that are internal to the entity. However, policies may assume broader force when they become the means by which an implementing agency interprets existing statutes (e.g., PHS Policy). Principles are broader in their scope and intended application, and are accepted generalizations about a topic that are frequently endorsed by many and diverse organizations (e.g., the U.S. Government Principles). Procedures (often called "operating procedures" or "standard operating procedures") are typically detailed, step-by-step processes meant to ensure the consistent application of institutional practices. Establishing standard operating proce-

MUST, SHOULD, AND MAY

Must indicates actions that the Committee for the Update of the Guide considers imperative and mandatory duty or requirement for providing humane animal care and use. Should indicates a strong recommendation for achieving a goal; however, the Committee recognizes that individual circumstances might justify an alternative strategy. May indicates a suggestion to be considered.

The Guide is written in general terms so that its recommendations can be applied in diverse institutions and settings that produce or use animals for research, teaching, and testing. This approach requires that users, IACUCs, veterinarians, and producers apply professional judgment in making specific

2

Animal Care and Use Program

The proper care and use of laboratory animals in research, testing, teaching, and production (animal use) require scientific and professional judgment based on the animals' needs and their intended use. An animal care and use program (hereafter referred to as the Program) comprises all activities conducted by and at an institution that have a direct impact on the well-being of animals, including animal and veterinary care, policies and procedures, personnel and program management and oversight, occupational health and safety, institutional animal care and use committee (IACUC) functions, and animal facility design and management.

> Program: The activities conducted by and at an institution that have a direct impact on the well-being of animals, including animal and veterinary care, policies and procedures, personnel and program management and oversight, occupational health and safety, IACUC functions, and animal facility design and management.

This chapter defines the overall Program and key oversight responsibilities and provides guidelines to aid in developing an effective Program. Chapters 3, 4, and 5 cover the details of Program components: environment, housing, and management; veterinary care; and physical plant, respectively. Each institution should establish and provide sufficient resources for a Program that is managed in accord with the *Guide* and in compliance with applicable regulations, policies, and guidelines.

REGULATIONS, POLICIES, AND PRINCIPLES

The use of laboratory animals is governed by an interrelated, dynamic

of literature related to animal science and use of animals is constantly evolving, requiring Programs to remain current with the information and best practices.

PROGRAM MANAGEMENT

An effective Program requires clearly de ned roles that align respon- sibility with regulatory and management authority. US federal law creates a statutory basis for the institutional of cial (IO), the attending veterinarian (AV), and the institutional animal care and use committee (IACUC). The Guide endorses these concepts as important operating principles for all US and non-US animal care and use programs. Effective leadership in and col- laboration among these three components, which not only oversee but also support animal users, are necessary (Lowman 2008; Van Sluyters 2008). In addition, interactions with regulatory and funding agencies and accredita- tion organizations are an integral part of the Program.

As summarized here and discussed throughout the Guide, the primary oversight responsibilities in the Program rest with the IO, the AV, and the IACUC. Their roles t in a de ned organizational structure where the reporting relationships, authorities, and responsibilities of each are clearly de ned and transparent. Together they establish policies and procedures, ensure regulatory compliance, monitor Program performance, and support high-quality science and humane animal use. A Program that includes these elements and establishes a balance among them has the best chance of ef- ciently using resources while attaining the highest standards of animal well- being and scienti c quality (Bayne and Garnett 2008; Van S(y)-3(n)-, etsiS(-3(0)-3(0)-
e121s12121i12121s8r1e121i12121s12121i1212121i128r12128r1h12
erogt nd is
ees, o3(u)-3(r)rT(c)-3(i)-3(m)-1100lindttg and nsur18tg
e9 o319 P9 g11 o319 w1i

animal science and medicine are rapidly changing and evolving disciplines. The institution should provide opportunities and support for regular professional development and continuing education to ensure both that professional staff are knowledgeable about the latest practices and procedures and that laboratory animals receive high-quality care (Colby et al. 2007).

Animal Care Personnel Personnel caring for animals should be appropriately trained (see Appendix A, Education), and the institution should provide for formal and/or on-the-job training to facilitate effective implementation of the Program and the humane care and use of animals. Staff should receive training and/or have the experience to complete the tasks for which they are responsible. According to the Program scope, personnel with expertise in various disciplines (e.g., animal husbandry, administration, veterinary medical technology) may be required.

There are a number of options for training animal care personnel and technicians (Pritt and Duffee 2007). Many colleges have accredited programs in veterinary technology (AVMA 2010); most are 2-year programs that award Associate of Science degrees, some are 4-year programs that award Bachelor of Science degrees. Nondegree training, via certification

used (Conarello and Shepard 2007). Training should be tailored to the particular needs of research groups; however, all research groups should receive training in animal care and use legislation, IACUC function, ethics of animal use and the concepts of the Three Rs, methods for reporting concerns about animal use, occupational health and safety issues pertaining to animal use, animal handling, aseptic surgical technique, anesthesia and analgesia, euthanasia, and other subjects, as required by statute. Continuing education programs should be offered to reinforce training and provide updates that re ect changes in technology, legislation, and other relevant areas. Frequency of training opportunities should ensure that all animal

there is a regulatory requirement for such a committee. Operational and day-to-day responsibility for safety in the workplace resides with the laboratory or facility supervisor (e.g., principal investigator, facility director, or a staff veterinarian) and depends on safe work practices by all employees.

Control and Prevention StrategiesA comprehensive OHSP should include a hierarchy of control and prevention strategies that begins with the identification of hazards and the assessment of risk associated with those hazards. Managing risk involves the following steps: rst, the appropriate design and operation of facilities and use of appropriate safety equipment (engineering controls); second, the development of processes and standard operating procedures (SOPs; administrative controls); and nally, the provision of appropriate personal protective equipment (PPE) for employees. Special safety equipment should be used in combination with appropriate management and safety practices (NIH 2002; OSHA 1998a,b). Managing risk using these strategies requires that personnel be trained, maintain good personal hygiene, be knowledgeable about the hazards in their work environment, understand the proper selection and use of equipment, follow established procedures, and use the PPE provided.

Hazard Identi cation and Risk AssessmentThe institutional OHSP should identify potential hazards in the work environment and conduct a critical assessment of the associated risks. An effective OHSP ensures that the risks associated with the experimental use of animals are identi ed and reduced to mttttto

severity or seriousness of the hazard); the exposure intensity, duration, and frequency (prevalence of the hazard); to some extent, the susceptibility (e.g., immune status) of the personnel; and the history of occupational illness and injury in the particular workplace (Newcomer 2002; NRC 1997). Ongoing identi cation and evaluation of hazards call for periodic inspections and reporting of potential hazardous conditions or “near miss” incidents.

Facilities, Equipment, and Monitoring The facilities required to support the OHSP will vary depending on the scope and activities of the Program. Their design should preferentially use engineering controls and equipment to minimize exposure to anticipated hazards (also see Chapter 5). Because a high standard of personal cleanliness is essential, changing, washing, and showering facilities and supplies appropriate to the Program should be available.

Where biologic agents are used, the Centers for Disease Control and Prevention (CDC) and National Institutes of Health (NIH) publication Biosafety in Microbiological and Biomedical Laboratories (BMBL; DHHS 2009) and the USDA standards (USDA 2002) should be consulted for appropriate facility design and safety procedures. These design and safety features are based on the level of risk posed by the agents used. Special facilities and safety equipment may be needed to protect the animal care and investi gative staff, other occupants of the facility, the public, animals, and the environment from exposure to hazardous biologic, chemical, and physical agents used in animal experimentation (DHHS 2009; Frasier and Talka 2005; NIH 2002). When necessary, these facilities should be separated from other animal housing and support areas, research and clinical laboratories, and patient care facilities. They should be appropriately identi ed and access to them limited to authorized personnel.

Facilities, equipment, and procedures should also be designed, selected, and developed to reduce the possibility of physical injury or health risk to personnel (NIOSH 1997a,b). Engineering controls and equipment that address the risk of ergonomic injury in activities such as the lifting of heavy equipment or animals should be considered (AVMA 2008). Those are also frequently used to limit or control personnel exposure to animal allergens (Harrison 2001; Huerkamp et al. 2009). The potential for repetitive motion injuries in animal facilities (e.g., maintenance of large rodent populations and other husbandry activities) should also be assessed.

The selection of appropriate animal housing systems requires profes sional knowledge and judgment and depends on the nature of the hazards in question, the types of animals used, the limitations or capabilities of the facilities, and the design of the experiments. Experimental animals should be housed so that possibly contaminated food and bedding, feces, and urine

Institutions should have written policies and procedures governing experimentation with hazardous biologic, chemical, and physical agents. An oversight process (such as the use of a safety committee) should be developed to involve persons who are knowledgeable in the evaluation and safe use of hazardous materials or procedures and should include review of the procedures and facilities to be used for speci c safety concerns. Formal safety programs should be established to assess hazards, determine the safeguards needed for their control, and ensure that staff have the necessary training and skills and that facilities are adequate for the safe conduct of the research. Technical support should be provided to monitor and ensure compliance with institutional safety policies. A collaborative approach involving the investigator and research team, attending veterinarian, animal care technician, and occupational health and safety professionals may enhance compliance.

The BMBL (DHHS 2009) and NRC (1997) recommend practices and procedures, safety equipment, and facility requirements for working with hazardous biologic agents and materials. Facilities that handle agents of unknown risk should consult with appropriate CDC personnel about hazard control and medical surveillance. The use of highly pathogenic "select agents and toxins" in research requires that institutions develop a program and procedures for procuring, maintaining, and disposing of these agents (CFR 1998, 2002a,b; NRC 2004; PL 107-56; PL 107-188; Richmond et al. 2003). The use of immunode cient or genetically modi ed animals (GMAs) susceptible to or shedding human pathogens, the use of human tissues and cell lines, or any infectious disease model can lead to an increased risk to the health and safety of personnel working with the animals (Lassnig et al. 2005; NIH 2002).

Hazardous agents should be contained in the study environment, for example through the use of air ow control during the handling and administering of hazardous agents, necropsies on contaminated animals (CDC and NIH 2000), and work with chemical hazards (Thomann 2003). Waste anesthetic gases should be scavenged to limit exposure.

Personal Protection While engineering and administrative controls are the

as gloves, arm protectors, suitable face masks, face shields, and goggles (NRC 2003a). Hearing protection should be available in high-noise areas (OSHA 1998c). Personnel working in areas where they might be exposed to contaminated airborne particulate material or vapors should have suitable respiratory protection (Fechter 1995; McCullough 2000; OSHA 1998d), with respirator fit testing and training in the proper use and maintenance of the respirator (OSHA 1998d; Sargent and Gallo 2003).

Medical Evaluation and Preventive Medicine for Personnel Development and implementation of a program of medical evaluation and preventive medicine should involve input from trained health professionals, such as occupational health physicians and nurses. Confidentiality and other medical and legal factors must be considered in the context of appropriate federal, state, and local regulations (e.g., PL 104-191).

A preemployment health evaluation and/or a health history evaluation before work assignment is advisable to assess potential risks for individual employees. Periodic medical evaluations are advisable for personnel in specific risk categories. For example, personnel required to use respiratory protection may also require medical evaluation to ensure that they

employee(s), unless such concerns are reported anonymously; corrective actions if deemed necessary; and a report to the IO of the issue, ndings, and actions taken. Reported concerns and any corrective actions taken should be documented.

Mechanisms for reporting concerns should be posted in prominent locations in the facility and on applicable institutional website(s) with instructions on how to report the concern and to whom. Multiple points of contact, including senior management, the IO, IACUC Chair, and AV, are recommended. The process should include a mechanism for anonymity, compliance with applicable whistleblower policies, nondiscrimination against the concerned/reporting party, and protection from reprisals.

Training and regular communication with employees (including personnel such as custodial, maintenance, and administrative staff, who are farther removed from the animal use) about the institution's animal use activities may reduce potential concerns.

PROGRAM OVERSIGHT

The Role of the IACUC

IACUC Constitution and Function

The responsibility of the IACUC is to oversee and routinely evaluate the Program. It is the institution's responsibility to provide suitable orientation, background materials, access to appropriate resources, and, if necessary, speci c training to assist IACUC members in understanding their roles and responsibilities and evaluating issues brought before the committee.

Committee membership includes the following:

- a Doctor of Veterinary Medicine either certi ed (e.g., by ACLAM, ECLAM, JCLAM, KCLAM) or with training and experience in laboratory animal science and medicine or in the use of the species at the institution
- at least one practicing scientist experienced in research involving animals
- at least one member from a nonscienti c background, drawn from inside or outside the institution
- at least one public member to represent general community inter-

3(a)-3(l)--3(g)-3(h)ID 2118 h3(e)-3(r)]TJ EMnd4(r)]TJ EMnceinng an1-3(e)-3(

compensation for participation and ancillary expenses (e.g., meals, parking, travel), but the amount should be sufficiently modest that it does not become a substantial source of income and thus risk compromising the member's association with the community and public at large.

For large institutions with many administrative units or departments, no more than three voting members should be associated with a single administrative unit (USDA 1985). The size of the institution and the nature and extent of the Program will determine the number of members of the committee and their terms of appointment. Institutions with broad research programs may need to choose scientists from a number of disciplines and experience to properly evaluate animal use protocols.

The committee is responsible for oversight and evaluation of the entire Program and its components as described in other sections of the Guide. Its oversight functions include review and approval of proposed animal use (protocol review) and of proposed significant changes to animal use; regular inspection of facilities and animal use areas; regular review of the Program; ongoing assessment of animal care and use; and establishment of a mechanism for receipt and review of concerns involving the care and use of animals at the institution. The committee must meet as often as necessary to fulfill its responsibilities, and records of committee meetings and results of deliberations should be maintained. Program review and facilities inspections should occur at least annually or more often as required (e.g., by the Animal Welfare Act and PHS Policy). After review and inspection, a written report (including any minority views) should be provided to the IO about the status of the Program.

Protocol Review

The animal use protocol is a detailed description of the proposed use of laboratory animals. The following topics should be considered in the preparation of the protocol by the researcher and its review by the IACUC:

- rationale and purpose of the proposed use of animals
- a clear and concise sequential description of the procedures involving the use of animals that is easily understood by all members of the committee
- availability or appropriateness of the use of less invasive procedures, other species, isolated organ preparation, cell or tissue culture, or computer simulation (see Appendix A, Alternatives)
- justification of the species and number of animals proposed; whenever possible, the number of animals and experimental group sizes should be statistically justified (e.g., provision of a power analysis; see Appendix A, Experimental Design and Statistics)

- unnecessary duplication of experiments

Special Considerations for IACUC Review

Certain animal use protocols include procedures or approaches that require special consideration during the IACUC review process due to their potential for unrelieved pain or distress or other animal welfare concerns. The topics below are some of the most common requiring special IACUC consideration. For these and other areas the IACUC is obliged to weigh the objectives of the study against potential animal welfare concerns. By considering opportunities for re nement, the use of appropriate nonanimal alternatives, and the use of fewer animals, both the institution and the principal investigator (PI) can begin to address their shared obligations for humane animal care and use.

Experimental and Humane EndpointsThe experimental endpoint of a study occurs when the scienti c aims and objectives have been reached. The humane endpoint is the point at which pain or distress in an experimental animal is prevented, terminated, or relieved. The use of humane endpoints contributes to re nement by providing an alternative to experimental end-

cbj a(o)-3(n)-3(t)-3(l)-3(t)-3((o)-3a)-3(le)-3(r)-3(o)110(v)-3(e)-3(nx)-3krdro())-3Werl nd2

humane endpoint, and the response required upon reaching the humane endpoint. An understanding of preemptive euthanasia (Toth 2000), behavioral or physiologic de nitions of the moribund state (ibid.), and the use of study-speci c animal assessment records (Morton 2000; Paster et al. 2009) can aid the PI and IACUC when considering or developing proposed endpoints. When novel studies are proposed or information for an alternative endpoint is lacking, the use of pilot studies is an effective method for identifying and de ning humane endpoints and reaching consensus among the PI, IACUC, and veterinarian. A system for communication with the IACUC should be in place both during and after such studies. Numerous publications address speci c proposals for the application and use of humane endpoints (e.g., CCAC 1998; ILAR 2000; OECD 1999; Toth 1997; UKCCCR 1997).

Unexpected Outcomes Fundamental to scienti c inquiry is the investiga-tion of novel experimental variables. Because of the potential for unex-pected outcomes that may affect animal well-being when --217f-3(a)-3(l)-3( )-h3(-

- Veterinary care must be provided if lesions or illnesses associated with restraint are observed. The presence of lesions, illness, or severe behavioral change often necessitates the temporary or permanent removal of the animal from restraint.
- The purpose of the restraint and its duration should be clearly explained to personnel involved with the study.

Multiple Survival Surgical Procedures Surgical procedures in the laboratory setting may be categorized as major or minor (USDA 1985). Whether a procedure is major or minor should be evaluated on a case-by-case basis, as determined by the veterinarian and IACUC (NRC 2003b; Silverman et al. 2007; for additional discussion see Chapter 4, Surgical Procedures).

Regardless of classification, multiple surgical procedures on a single animal should be evaluated to determine their impact on the animal's well-being. Multiple major surgical procedures on a single animal are acceptable only if they are (1) included in and essential components of a single research project or protocol, (2) scientifically justified by the investigator, or (3) necessary for clinical reasons. Conservation of scarce animal resources may justify the conduct of multiple major surgeries on a single animal, but the application of such a practice on a single animal used in separate protocols is discouraged and should be reviewed critically by the IACUC. When applicable, the IO must submit a reQsw40 [(()-3()-3(O)-3( )-112(m)-3(u)-3(Dtg

least restriction necessary to achieve the scientific objective while maintaining animal well-being.

The development of animal protocols that involve the use of food or fluid regulation requires the evaluation of three factors: the necessary level of regulation, potential adverse consequences of regulation, and methods for assessing the health and well-being of the animals (NRC 2003b). In addition, the following factors influence the amount of food or fluid restriction that can be safely used in a specific protocol: the species, strain, or stock, gender, and age of the animals; thermoregulatory demand; type of housing; time of feeding, nutritive value, and fiber content of the diet (Heiderstadt et al. 2000; Rowland 2007); and prior experimental manipulation. The degree of food or fluid restriction necessary for consistent behavioral performance is influenced by the difficulty of the task, the individual animal, the motivation required of the animal, and the effectiveness of animal training for a specific protocol-related task.

The animals should be closely monitored to ensure that food and fluid intake meets their nutritional needs (Toth and Gardiner 2000). Body weights should be recorded at least weekly and more often for animals requiring greater restrictions (NRC 2003b). Written records should be maintained for each animal to document daily food and fluid consumption, hydration status, and any behavioral and clinical changes used as criteria for temporary or permanent removal of an animal from a protocol (Morton 2000; NRC 2003b). In the case of conditioned-response research protocols, use of a highly preferred food or fluid as positive reinforcement, instead of restriction, is recommended. Caloric restriction, as a husbandry technique and means of weight control, is discussed in Chapter 3.

Use of Non-Pharmaceutical-Grade Chemicals and Other Substances The

Field Investigations Investigations may involve the observation or use of nondomesticated vertebrate species under eld conditions. Many eld investigations require international, federal, state, and/or local permits, which may call for an evaluation of the scienti c merit of the proposed study and a determination of the potential impact on the population or species to be studied.

Additionally, occupational health and safety issues, including zoonoses, should be reviewed by the institution's health and safety committee or of ce, with assurances to the IACUC that the eld study does not compromise the health and safety of either animals or persons in the eld. Principal investigators conducting eld research should be knowledgeable about relevant zoonotic diseases, associated safety issues, and any laws or regulations that apply. Exceptions to the above should be clearly de ned and evaluated by the IACUC.

In preparing the design of a eld study, investigators are encouraged to consult with relevant professional societies and available guidelines (see Appendix A). Veterinary input may be needed for projects involving capture, individual identi cation, sedation, anesthesia, surgery, recovery, holding, transportation, release, or euthanasia. Issues associated with these activities are similar if not identical to those for species maintained and used in the laboratory. When species are removed from the wild, the protocol should include plans for either a return to their habitat or their nal disposition, as appropriate.

The Guide does not purport to be a compendium of all information regarding eld biology and methods used in wildlife investigations, but the basic principles of humane care and use apply to animals living under natural conditions. IACUCs engaged in the review of eld studies are encouraged to consult with a quali ed wildlife biologist.

Agricultural Animals The use of agricultural animals in research is subject to the same ethical considerations as for other animals in research, although it is often categorized as either biomedical or agricultural because of government regulations and policies, institutional policies, administrative structure, funding sources, and/or user goals (Stricklin et al. 1990). This categorization has led to a dual system with different criteria for evaluating protocols and standards of housing and care for animals of the same species on the basis of stated biomedical or agricultural research objectives (Stricklin and Mench 1994). With some studies, differences in research goals may lead to a clear distinction between biomedical and agricultural research. For example, animal models of human diseases, organ transplantation, and major surgery are considered biomedical u fd

disease studies, administrators, regulators, and IACUCs face a dilemma in deciding how to handle such studies (Stricklin et al. 1990). Decisions on categorizing research uses of agricultural animals and de ning standards for their care and use should be made by the IACUC based on both the researcher's goals and concern for animal well-being. Regardless of the category of research, institutions are expected to provide oversight of all research animals and ensure that pain and distress are minimized.

The protocol, rather than the category of research, should determine the setting (farm or laboratory). Housing systems for agricultural animals used in biomedical research may or may not differ from those used in agricultural research; animals used in either type of research can be housed in cages, stalls, paddocks, or pastures (Tillman 1994). Some agricultural studies need uniform conditions to minimize environmental variability, and some biomedical studies are conducted in farm settings. Agricultural research often necessitates that animals be managed according to contemporary farm production practices (Stricklin and Mench 1994), and natural environmental conditions might be desirable for agricultural research, whereas control of environmental conditions to minimize variation might be desirable in biomedical research (Tillman 1994).

The Guide applies to agricultural animals used in biomedical research, including those maintained in typical farm settings. For animals maintained in a farm setting, the*Guide for the Care and Use of Agricultural Animals in Research and Teaching*(FASS 2010) is a useful resource. Information about environmental enrichment, transport, and handling may be helpful in both agricultural and biomedical research settings. Additional information about facilities and management of farm animals in an agricultural setting is available from the Midwest Plan Service (1987) and from agricultural engineers or animal science experts.

Postapproval Monitoring

Continuing IACUC oversight of animal activities is required by federal laws, regulations, and policies. A variety of mechanisms can be used to facilitate ongoing protocol assessment and regulatory compliance. Postapproval monitoring (PAM) is considered here in the broadest sense, consist

ments. The IACUC, veterinary, animal care, and compliance staff may all conduct PAM, which may also serve as an educational tool.

Continuing protocol review typically consists of an annual update or review as well as the triennial review required by the PHS. The depth of such reviews varies from a cursory update to a full committee review of the entire protocol. Some institutions use the annual review as an opportunity for the investigator to submit proposed amendments for future procedures, to provide a description of any adverse or unanticipated events, and to provide updates on work progress. For the triennial review, many institutions require a complete new protocol submission and may request a progress report on the use of animals during the previous 3 years.

Both the Health Research Extension Act and the AWA require the IACUC to inspect animal care and use facilities, including sites used for animal surgeries, every 6 months. As part of a formal PAM program some institutions combine inspection of animal study sites with concurrent review of animal protocols. Based on risks to animals and their handlers, other study areas may require more or less frequent inspections. Examples of effective monitoring strategies include

- examination of surgical areas, including anesthetic equipment, use of appropriate aseptic technique, and handling and use of controlled substances
- review of protocol-related health and safety issues
- review of anesthetic and surgical records
- regular review of adverse or unexpected experimental outcomes affecting the animals
- observation of laboratory practices and procedures and comparison with approved protocols.

Institutions may also consider the use of veterinary staff and/or animal health technicians to observe increased risk procedures for adverse events (e.g., novel survival surgeries, pain studies, tumor growth studies) and report their findings for review by the IACUC. The level of formality and intensity of PAM should be tailored to institutional size and complexity, and in all cases should support a culture of care focusing on the animals' well-being (Klein and Bayne 2007). Regardless of the methods used or who conducts and coordinates the monitoring, PAM programs are more likely to succeed when the institution encourages an educational partnership with investigators (Banks and Norton 2008; Collins 2008; Dale 2008; Lowman 2008; Plante and James 2008; Van Sluyters 2008).

DISASTER PLANNING AND EMERGENCY PREPAREDNESS

Animal facilities may be subject to unexpected conditions that result in the catastrophic failure of critical systems or signi cant personnel absentee ism, or other unexpected events that severely compromise ongoing animal care and well-being (ILAR 2010). Facilities must therefore have a disaster plan. The plan should de ne the actions necessary to prevent animal pain, distress, and deaths due to loss of systems such as those that control venti lation, cooling, heating, or provision of potable water. If possible the plan should describe how the facility will preserve animals that are necessary for critical research activities or are irreplaceable. Knowledge of the geo graphic locale may provide guidance as to the probability of a particular type of disaster.

Disaster plans should be established in conjunction with the respon sible investigator(s), taking into consideration both the priorities for triaging animal populations and the institutional needs and resources. Animals that cannot be relocated or protected from the consequences of the disaster must be humanely euthanized. The disaster plan should identify essential person nel who should be trained in advance in its implementation. Efforts should be taken to ensure personnel safety and provide access to essential person nel during or immediately after a disaster. Such plans should be approved by the institution and be part of the overall institutional disaster response plan that is coordinated by the IO or another senior-level administrator. Law enforcement and emergency personnel should be provided with a copy of the plan for comment and integration into broader, areawide planning (Vogelweid 1998).

REFERENCES

AAALAC [Association for Assessment and Accreditation of Laboratory Animal Care] Interna tional. 2003. Who's responsible for offsite animals? Connection Spring:6-11, 13. Available at www.aaalac.org/publications.

ACLAM [American College of Laboratory Animal Medicine]. 1996. Adequate Veterinary Care. Available at www.aclam.org/education/guidelines/position_adequatecare.html; accessed May 10, 2010.

Anderson LC. 2007. Institutional and IACUC responsibilities for animal care and use education and training programs. ILAR J 48:90-95.

AVMA [American Veterinary Medical Association]. 2008. Introduction to Ergonomics Guide lines for Veterinary Practice. April. Available at www.avma.org/issues/policy/ergonomics. asp; accessed May 10, 2010.

AVMA. 2010. Programs accredited by the AVMA Committee on Veterinary Technician Education and Activities (CVTEA). Available at www.avma.org/education/cvea/vettech_ programs/vettech_programs.asp; accessed January 4, 2010.

Banks RE, Norton JN. 2008. A sample postapproval monitoring program in academia. ILAR J 49:402-418.

Bayne KA, Garnett NL. 2008. Mitigating risk, facilitating research. ILAR J 49:369-371.

Boissy A, Manteuffel G, Jensen MB, Moe RO, Spruijt B, Keeling L, Winckler C, Forkman B, Dimitrov I, Langbein J, Bakken M, Veissier I, Aubert A. 2007. Assesment of positive emotions in animals to improve their welfare. Physiol Behav 92:375-397.

DHHS [Department of Health and Human Services]. 2009. Biosafety in Microbiological and Biomedical Laboratories, 5th ed. Chosewood LC, Wilson DE, eds. Washington: Government Printing Office. Available at http://www.cdc.gov/biosafety/publications/bmbl5/index.htm; accessed July 30, 2010.

FASS [Federation of Animal Science Societies]. 2010. Guide for the Care and Use of Agricultural Animals in Research and Teaching, 3rd ed. Champlain, IL: FASS.

Mann MD, Prentice ED. 2004. Should IACUCs review scientific merit of animal research projects? Lab Anim (NY) 33:26-31.

McCullough NV. 2000. Personal respiratory protection. In: Fleming DO, Hunt DL, eds. Biological Safety Principles and Practices. Washington: ASM Press. p 383-404.

Meunier LD. 2006. Selection, acclimation, training and preparation of dogs for the research setting. ILAR J 47:326-347.

Midwest Plan Service. 1987. Structures and Environment Handbook, 11th ed. rev. Ames: Midwest Plan Service, Iowa State University.

Miller G. 2007. Science and the public: Animal extremists get personal. Science 318:1856-1858.

Moorehead RA, Sanchez OH, Baldwin RM, Khokha R. 2003. Transgenic overexpression of IGF-II induces spontaneous lung tumors: A model for human lung adenocarcinoma. Oncogene 22:853-857.

Morton DB. 2000. A systematic approach for establishing humane endpoints. ILAR J 41:80-86.

Morton WR, Knitter GH, Smith PV, Susor TG, Schmitt K. 1987. Alternatives to chronic restraint of nonhuman primates. JAVMA 191:1282-1286.

Stokes WS. 2000. Reducing unrelieved pain and distress in laboratory animals using humane endpoints. ILAR J 41:59-61.

Stokes WS. 2002. Humane endpoints for laboratory animals used in regulatory testing. ILAR J 43:S31-S38.

Stricklin WR, Mench JA. 1994. Oversight of the use of agricultural animals in university teaching and research. ILAR News 36:9-14.

Stricklin WR, Purcell D, Mench JA. 1990. Farm animals in agricultural and biomedical research. In: The Well-Being of Agricultural Animals in Biomedical and Agricultural Research: Proceedings from a SCAW-Sponsored Conference, September 6-7. Washington: Scientists Center for Animal Welfare. p 1-4.

Thomann WR. 2003. Chemical safety in animal care, use, and research. ILAR J 44:13-19.

Thulin H, Bjorkdahl M, Karlsson AS, Renstrom A. 2002. Reduction of exposure to laboratory animal allergens in a research laboratory. Ann Occup Hyg 46:61-68.

Tillman P. 1994. Integrating agricultural and biomedical research policies: Conflicts and opportunities. ILAR News 36:29-35.

Toth LA. 1997. The moribund state as an experimental endpoint. Contemp Top Lab Anim Sci 36:44-48.

Toth LA. 2000. Defining the moribund condition as an experimental endpoint for animal research. ILAR J 41:72-79.

Toth LA, Gardiner TW. 2000. Food and water restriction protocols: Physiological and behavioral considerations. Contemp Top Lab Anim Sci 39:9-17.

UKCCCR [United Kingdom Coordinating Committee on Cancer Research]. 1997. Guidelines for the Welfare of Animals in Experimental Neoplasia, 2nd ed. London: UKCCCR.

USC [United States Code]. Title 42, Chapter 6a, Subchapter III, Part H, Section 289d: Animals in Research. Available at http://uscode.house.gov/download/pls/42CGA.txt.

USDA [US Department of Agriculture]. 1985. 9 CFR 1A. (Title 9, Chapter 1, Subchapter A): Animal Welfare. Available at http://ecfr.gpoaccess.gov/cgi/t/text/text-idx?sid=8314313bd7adf2c9f1964e2d82a88d92andc=ecfrandtpl=/ecfrbrowse/Title09/9cfrv1_02.tpl; accessed January 14, 2010.

USDA. 1997a. APHIS Policy #14, “Multiple Survival Surgery: Single vs. Multiple Procedures” (April 14). Available at www.aphis.usda.gov/animal_welfare/downloads/policy/policy14.pdf; accessed January 4, 2010.

USDA. 1997b. APHIS Policy #3, “Veterinary Care” (July 17). Available at www.aphis.usda.gov/animal_welfare/downloads/policy/policy3.pdf; accessed January 9, 2010.

USDA. 2002. Facilities Design Standards. Manual 242.1. Available at www.afm.ars.usda.gov/ppweb/PDF/242-01M.pdf; accessed May 10, 2010.65(a)-3(t)-3((2)-3(0)-3)-31(a)-3(c)-3(c)-3(e)-3(

reptiles and certain amphibian species, are terrestrial. Personnel working with aquatic animals should be familiar with management implications, e.g., the importance of providing appropriate temperature ranges for basic physiologic function.

TERRESTRIAL ANIMALS

Terrestrial Environment

Microenvironment and Macroenvironment

The microenvironment of a terrestrial animal is the physical environment immediately surrounding it; that is, the primary enclosure such as the cage, pen, or stall. It contains all the resources with which the animals come directly in contact and also provides the limits of the animals' immediate environment. The microenvironment is characterized by many factors, including illumination, noise, vibration, temperature, humidity, and gaseous and particulate composition of the air. The physical environment of the secondary enclosure, such as a room, a barn, or an outdoor habitat, constitutes the macroenvironment.

Microenvironment: The immediate physical environment surrounding the animal (i.e., the environment in the primary enclosure such as the cage, pen, or stall).

Macroenvironment: The physical environment of the second factors, int

Temperature and Humidity

environmental relative humidity may be of greater importance for animals housed in a primary enclosure in which the environmental conditions differ greatly from those of the macroenvironment (e.g., in static lter-top [isola tor] cages).

Some species may require conditions with high relative humidity (e.g.,

and should therefore be considered. The use of computer modeling for assessing those factors in relation to heat loading, air diffusion patterns, and particulate movement may be helpful in optimizing ventilation of micro- and macroenvironments (Hughes and Reynolds 1995).

Direct exposure of animals to air moving at high velocity (drafts) should be avoided as the speed of air to which animals are exposed affects the rate at which heat and moisture are removed from an animal. For example, air at 20°C moving at 60 linear feet per minute (18.3 m/min) has a cooling effect of approximately 7°C (Weihe 1971). Drafts can be particularly problematic for neonatal homeotherms (which may be hairless and have poorly devel oped mechanisms for thermoregulatory control), for mutants lacking fur, and for semiaquatic amphibians that can desiccate.

Provision of 10 to 15 fresh air changes per hour in animal housing rooms is an acceptable guideline to maintain macroenvironmental air qual ity by constant volume systems and may also ensure microenvironmental air quality. Although this range is effective in many animal housing settings, it does not take into account the range of possible heat loads; the species, size, and number of animals involved; the type of primary enclosure and bedding; the frequency of cage changing; the room dimensions; or the ef ciency of air distribution both in the macroenvironment and between the macro- and microenvironments. In some situations, the use of such a broad guideline might overventilate a macroenvironment containing few animals, thereby wasting energy, or underventilate a microenvironment containing many animals, allowing heat, moisture, and pollutants to accumulate.

Modern heating, ventilation, and air conditioning (HVAC) systems (e.g., variable air volume, or VAV, systems) allow v1(u)-3(n)-3(d)-3(e)-3(r)-3(ν)-3(e)-3(n)-3

able to exhaust these systems directly into the building's exhaust system to reduce heat load and macroenvironmental contamination.

well as the duration of the animal's current and prior exposure to light, and the animal's pigmentation, circadian rhythm, body temperature, hormonal status, age, species, sex, and stock or strain (Brainard 1989; Duncan and O'Steen 1985; O'Steen 1980; Saltarelli and Coppola 1979; Semple-Rowland and Dawson 1987; Wax 1977). More recent studies in rodents and primates have shown the importance of intrinsically photosensitive retinal ganglion cells (distinct from rods and cones) for neuroendocrine, circadian, and neurobehavioral regulation (Berson et al. 2002; Hanifin and Brainard 2007). These cells can respond to light wavelengths that may differ from other photoreceptors and may influence the type of lighting, light intensity, and wavelength selected for certain types of research.

In general, lighting should be diffused throughout an animal holding area and provide sufficient illumination for the animals' well-being while permitting good housekeeping practices, adequate animal inspection

midcage (NASA 1988). Rats and mice generally prefer cages with low light intensity (Blom et al. 1996), and albino rats prefer areas with a light intensity of less than 25 lux (Schlingmann et al. 1993a). Young mice prefer much lower illumination than adults (Wax 1977). For animals that have been shown to be susceptible to phototoxic retinopathy, light should be between 130 and 325 lux in the room at cage level.

Light intensity decreases with the square of the distance from its source. Thus the location of a cage on a rack affects the intensity of light to which the animals within are exposed. Light intensity may differ as much as 80-fold in transparent cages from the top to the bottom of a rack, and differences up to 20-fold have been recorded within a cage (Schlingmann et al. 1993a,b). Management practices, such as rotating cage position relative to the light source (Greenman et al. 1982) or providing animals with ways to control their own light exposure by behavioral means (e.g., nesting or bedding material adequate for tunneling), can reduce inappropriate light stimulation. Variable-intensity lights are often used to accommodate the needs of research protocols, certain animal species, and energy conservation. However, such a system should also provide for the observation and care of the animals. Caution should be exercised as increases in daytime room illumination for maintenance purposes have been shown to change photoreceptor physiology and can alter circadian regulation (NRC 1996; Reme et al. 1991; Terman et al. 1991).

Noise and Vibration

Noise produced by animals and animal care activities is inherent in the

physical, physiologic, and behavioral needs. Environments that fail to meet the animals' needs may result in abnormal brain development, physiologic dysfunction, and behavioral disorders (Garner 2005; van Praag et al. 2000; Würbel 2001) that may compromise both animal well-being and scienti c validity. The primary enclosure or space may need to be enriched to prevent such effects (see also section on Environmental Enrichment).

An appropriate housing space or enclosure should also account for the animals' social needs. Social animals should be housed in stable pairs or groups of compatible individuals unless they must be housed alone for experimental reasons or because of social incompatibility (see also section on Behavioral and Social Management). Structural adjustments are fre quently required for social housing (e.g., perches, visual barriers, refuges), and important resources (e.g., food, water, and shelter) should be provided in such a way that they cannot be monopolized by dominant animals (see also section on Environmental Enrichment).

The primary enclosure should provide a secure environment that does not permit animal escape and should be made of durable, nontoxic materi als that resist corrosion, withstand the rigors of cleaning and regular han dling, and are not detrimental to the health and research use of the animals. The enclosure should be designed and manufactured to prevent accidental entrapment of animals or their appendages and should be free of sharp edges or projections that could cause injury to the animals or personnel. It should have smooth, impervious surfaces with minimal ledges, angles, corners, and overlapping surfaces so that accumulation of dirt, debris, and moisture is minimized and cleaning and disinfecting are not impaired. All enclosures should be kept in good repair to prevent escape of or injury to animals, promote physical comfort, and facilitate sanitation and servic ing. Rusting or oxidized equipment, which threatens the health or safety of animals, needs to be repaired or replaced. Less durable materials, such as wood, may be appropriate in select situations, such as outdoor corrals, perches, climbing structures, resting areas, and perimeter fences for primary enclosures. Wooden items may need to be replaced periodically because of damage or dif culties with sanitation. Painting or sealing wood surfaces with nontoxic materials may improve durability in many instances.

Flooring should be solid, perforated, or slatted with a slip-resistant sur face. In the case of perforated or slatted oors, the holes and slats should have smooth edges. Their size and spacing need to be commensurate with the size of the housed animal to minimize injury and the development of foot lesions. If wire-mesh ooring is used, a solid resting area may be bene cial, as this oor type can induce foot lesions in rodents and rabbits (Drescher 1993; Fullerton and Gilliatt 1967; Rommers and Meijerhof 1996). The size and weight of the animal as well as the duration of housing on wire-mesh oors may also play a role in the development of this condi-

exercise, manipulative activities, and cognitive challenges according to species-speci c characteristics (NRC 1998a; Young 2003). Examples of enrichment include structural additions such as perches and visual barriers for nonhuman primates (Novak et al. 2007); elevated shelves for cats (Overall and Dyer 2005; van den Bos and de Cock Buning 1994) and rabbits (Stauffacher 1992); and shelters for guinea pigs (Baumans 2005), as well as manipulable resources such as novel objects and foraging devices for nonhuman primates; manipulable toys for nonhuman primates, dogs, cats, and swine; wooden chew sticks for some rodent species; and nesting material for mice (Gaskill et al. 2009; Hess et al. 2008; Hubrecht 1993; Lutz and Novak 2005; Olsson and Dahlborn 2002). Novelty of enrichment through rotation or replacement of items should be a consideration; how ever, changing animals' environment too frequently may be stressful.

Well-conceived enrichment provides animals with choices and a degree of control over their environment, which allows them to better cope with environmental stressors (Newberry 1995). For example, visual barriers allow nonhuman primates to avoid social con ict; elevated shelves for rabbits and shelters for rodents allow them to retreat in case of disturbances (Baumans 1997; Chmiel and Noonan 1996; Stauffacher 1992); and nesting material and deep bedding allow mice to control their temperature and avoid cold stress during resting and sleeping (Gaskill et al. 2009; Gordon 1993, 2004).

Not every item added to the animals' environment bene ts their well-being. For example, marbles are used as a stressor in mouse anxiety studies (De Boer and Koolhaas 2003), indicating that some items may be detrimen tal to well-being. For nonhuman primates, novel objects can increase the risk of disease transmission (Bayne et al. 1993); foraging devices can lead to increased body weight (Brent 1995); shavings can lead to allergies and skin rashes in some individuals; and some objects can result in injury from foreign material in the intestine (Hahn et al. 2000). In some strains of mice, cage dividers and shelters have induced overt aggression in groups of males, resulting in social stress and injury (e.g., Bergmann et al. 1994; Haemisch et al. 1994). Social stress was most likely to occur when resources were monopolized by dominant animals (Bergmann et al. 1994).

Enrichment programs should be reviewed by the IACUC, researchers, and veterinarian on a regular basis to ensure that they are bene cial to ani mal well-being and consistent with the goals of animal use. They should be updated as needed to ensure that they re ect current knowledge. Personnel responsible for animal care and husbandry should receive training in the behavioral biology of the species they work with to appropriately monitor the effects of enrichment as well as identify the development of adverse or abnormal behaviors.

Like other environmental factors (such as space, light, noise, tempera ture, and animal care procedures), enrichment affects animal phenotype

and may affect the experimental outcome. It should therefore be considered an independent variable and appropriately controlled.

Some scientists have raised concerns that environmental enrichment mc0-3(a-3(n)sa)-3,rii ii3-39 (i)12(i3-39T u(i)12(i3-33.i)-3(a)-3.i)-3-3e ietiana3(a)-3.3-M

changes when animals are first introduced to outdoor housing; training of animals to cooperate with veterinary and investigative personnel (e.g., to enter chutes or cages for restraint or transport); and adequate security via a perimeter fence or other means.

Naturalistic Environments

Areas such as pastures and islands may provide a suitable environment

TABLE 3.2 Recommended Minimum Space for Commonly Used Laboratory Rodents Housed in Groups*

Animals	Weight, g	Floor Area/Animal,[a] in.^{2} (cm^2)	Height,[b] in. (cm)	Comments
Mice in groups[c]	<10	6 (38.7)	5 (12.7)	Larger animals may require more space to meet the performance standards.
	Up to 15	8 (51.6)	5 (12.7)	
	Up to 25	12 (77.4)	5 (12.7)	
	>25	15 (96.7)	5 (12.7)	
Female + litter		51 (330) (recommended space for the housing group)	5 (12.7)	Other breeding configurations may require more space and will depend on considerations such as number of adults and litters, and size and age of litters.[d]
Rats in groups[c]	<100	17 (109.6)	7 (17.8)	Larger animals may require more space to meet the performance standards.
	Up to 200	23 (148.35)	7 (17.8)	
	Up to 300	29 (187.05)	7 (17.8)	
	Up to 400	40 (258.0)	7 (17.8)	
	Up to 500	60 (387.0)	7 (17.8)	
	>500	70 (451.5)	7 (17.8)	
Female + litter		124 (800) (recommended space for the housing group)	7 (17.8)	Other breeding configurations may require more space and will depend on considerations such as number of adults and litters, and size and age of litters.[d]
Hamsters[c]	<60	10 (64.5)	6 (15.2)	Larger animals may require more space to meet the performance standards.
	Up to 80	13 (83.8)	6 (15.2)	
	Up to 100	16 (103.2)	6 (15.2)	
	>100	19 (122.5)	6 (15.2)	
Guinea pigs[c]	Up to 350	60 (387.0)	7 (17.8)	Larger animals may require more space to meet the performance standards.
	>350	101 (651.5)	7 (17.8)	

*The interpretation of this table should take into consideration the performance indices described in the text beginning on page 55.

[a]Singly housed animals and s-3(r)-3(m)-3(a)-3(l)-3(l)-3(p)-3(c)-3(r)-3(o)-3(u h16ab <</MCID3(u h16abhnre8(t

animal may be required, while larger groups may be housed at slightly higher densities.

Studies have recently evaluated space needs and the effects of social housing, group size, and density (Andrade and Guimaraes 2003; Bartolomucci et al. 2002, 2003; Georgsson et al. 2001; Gonder and Laber 2007; Perez et al. 1997; A.L. Smith et al. 2004), age (Arakawa 2005; Davidson et al. 2007; Yildiz et al. 2007), and housing conditions (Gordon et al. 1998; Van Loo et al. 2004) for many different species and strains of rodents, and have reported varying effects on behavior (such as aggression) and experimental outcomes (Karolewicz and Paul 2001; Laber et al. 2008; McGlone et al. 2001; Rock et al. 1997; Smith et al. 2005; Van Loo et al. 2001). However, it is dif cult to compare these studies due to the study design and experimental variables that have been measured. For example, variables that may affect the animals' response to different cage sizes and housing densities include, but are not limited to, species, strain (and social behavior of the strain), phenotype, age, gender, quality of the space (e.g., vertical access), and structures placed in the cage. These issues remain complex and should be carefully considered when housing rodents.

Wolfensohn 2004). Group composition is critical and numerous species-speci c factors such as age, behavioral repertoire, sex, natural social orga nization, breeding requiremen3(z)-3(a) /V184(r)-3(e)-3()-3Mecituctnasc-herl84(r)-3(e)

TABLE 3.4 Recommended Minimum Space for Avian Species Housed in Pairs or Groups*

Animals	Weight,[a] kg	Floor area/animal,[b] ft^2 (m^2)	Height
Pigeons	—	0.8 (0.07)	Cage height should be
Quail	—	0.25 (0.023)	suf cient for the animals to
Chickens	<0.25	0.25 (0.023)	comfortably stand erect with
	Up to 0.5	0.50 (0.046)	their feet on the oor.
	Up to 1.5	1.00 (0.093)	
	Up to 3.0	2.00 (0.186)	
	>3.0[c]	3.00 (0.279)	

*The interpretation of this table should take into consideration the performance indices described in the text beginning on page 55.

[a]To convert kilograms to pounds, multiply by 2.2.
[b]Singly housed birds may require more space per animal than recommended for pair- or group-housed birds.
[c]Larger animals may require more space to meet performance standards (see text).

If it is necessary to house animals singly—for example, when justi ed for experimental purposes, for provision of veterinary care, or for incompat ible animals—this arrangement should be for the shortest duration possible. If single animals are housed in small enclosures, an opportunity for periodic release into larger enclosures with additional enrichment items should be considered, particularly for animals housed singly for extended periods of time. Singly housed animals may require more space per animal than recommended for pair- or group-housed animals, while larger groups may be housed at slightly higher densities. Because of the many physical and behavioral characteristics of nonhuman primate species and the many fac tors to consider when using these animals in a biomedical research setting, species-speci c plans for housing and management should be developed. Such plans should include strategies for environmental and psychological enrichment.

Agricultural Animals Table 3.6 lists recommended minimum space for agri cultural animals commonly used in a laboratory setting. As social animals, they should be housed in compatible pairs or larger groups of compatible animals. When animals exceed the weights in the table, more space is required. For larger animals (particularly swine) it is important that the con guration of the space allow the animals to turn around and move freely (Becker et al. 1989; Bracke et al. 2002). Food troughs and water devices should be provided in suf cient numbers to allow ready access for all ani mals. Singly housed animals may require more space than recommended in

TABLE 3.6 Recommended Minimum Space for Agricultural Animals*

Animals/Enclosure	Weight,[a] kg	Floor Area/Animal,[b] ft^2 (m^2)
Sheep and Goats		
1	<25	10.0 (0.9)
	Up to 50	15.0 (1.35)
	>50[c]	20.0 (1.8)
2-5	<25	8.5 (0.76)
	Up to 50	12.5 (1.12)
	>50[c]	17.0 (1.53)
>5	<25	7.5 (0.67)
	Up to 50	11.3 (1.02)
	>50[c]	15.0 (1.35)
Swine		
1	<15	8.0 (0.72)
	Up to 25	12.0 (1.08)
	Up to 50	15.0 (1.35)
	Up to 100	24.0 (2.16)
	Up to 200	48.0 (4.32)
	>200[c]	60.0 (5.4)
2-5	<25	6.0 (0.54)
	Up to 50	10.0 (0.9)
	Up to 100	20.0 (1.8)
	Up to 200	40.0 (3.6)
	>200	

>5	<75	18.0 (1.62)
	Up to 200	36.0 (3.24)
	Up to 350	54.0 (4.86)
	Up to 500	72.0 (6.48)
	Up to 650	93.0 (8.37)
	>650[c]	d108.0 (d9.72)
Horses	—	144.0 (12.96)
Ponies		
1-4	—	72.0 (6.48)
>4/Pen	≤200	60.0 (5.4)
	>200[c]	d72.0 (d6.48)

*The interpretation of this table should take into consideration the performance indices described in the text beginning on page 55.

[a]To convert kilograms to pounds, multiply by 2.2.
[b]Floor area con guration should be such that animals can turn around and move freely without touching food or water troughs, have ready access to food and water, and have suf cient space to comfortably rest away from areas soiled by urine and feces (see text).
[c]Larger animals may require more space to meet performance standards including suf cient space to turn around and move freely (see text).

the table to enable them to turn around and move freely without touching food or water troughs, have ready access to food and water, and have suf cient space to comfortably rest away from areas soiled by urine and feces.

Terrestrial Management

Behavioral and Social Management

Activity Animal Activity typically implies motor activity but also includes cognitive activity and social interaction. Animals' natural behavior and activity pro le should be considered during evaluation of suitable housing or behavioral assessment.

Animals maintained in a laboratory environment are generally restricted in their activities compared to free-ranging animals. Forced activity for reasons other than attempts to meet therapeutic or approved protocol objectives should be avoided. High levels of repetitive, unvarying behav ior (stereotypies, compulsive behaviors) may re ect disruptions of normal behavioral control mechanisms due to housing conditions or management practices (Garner 2005; NRC 1998a).

Dogs, cats, rabbits, and man_]TJ T* [(o)-3(b)871.8 -1.2 Td Mv4amTd Mv4

additional opportunities for activity by being walked on a leash, having access to a run, or being moved into areas for social contact, play, or expło ration (Wolff and Rupert 1991). Loa ng areas, exercise lots, and pastures are suitable for large farm animals, such as sheep, horses, and cattle.

Social Environment Appropriate social interactions among members of the

Institutions should urge feed vendors to periodically provide data from laboratory-based feed analyses for critical nutrients. The user should know the date of manufacture and other factors that affect the food's shelf life. Stale food or food transported and stored inappropriately can become deficient in nutrients. Upon receipt, bags of feed should be examined to ensure that they are intact and unstained to help ensure that their contents have not been potentially exposed to vermin, penetrated by liquids, or contaminated. Careful attention should be paid to quantities received in each shipment, and stock should be rotated so that the oldest food is used first.

Areas in which diets and diet ingredients are processed or stored should be kept clean and enclosed to prevent the entry of pests. Food stocks should be stored off the floor on pallets, racks, or carts in a manner that facilitates sanitation. Opened bags of food should be stored in vermin-proof containers to minimize contamination and to avoid the potential spread of pathogens. Exposure to elevated storage room temperatures, extremes in relative humidity, unsanitary conditions, and insects and other vermin hastens food deterioration. Storage of natural-ingredient diets at less than 21°C (70°F) and below 50% relative humidity is recommended. Precautions should be taken if perishable items—such as meats, fruits, and vegetables and some specialty diets (e.g., select medicated or high-fat diets)—are fed, because storage conditions may lead to variation in food quality.

Most natural-ingredient, dry laboratory animal diets stored properly can be used up to 6 months after manufacture. Nonstabilized vitamin C in manufactured feeds generally has a shelf life of only 3 months, but commonly used stabilized forms can extend the shelf life of feed. Refrigeration preserves nutritional quality and lengthens shelf life, but food storage time

ensure access to food for all animals, especially if feed is restricted as part of the protocol or management routine. Food storage containers should not be transferred between areas that pose different risks of contamination without appropriate treatment, and they should be cleaned and sanitized regularly.

Management of caloric intake is an accepted practice for long-term housing of some species, such as some rodents, rabbits, and nonhuman primates, and as an adjunct to some clinical, experimental, and surgical procedures (for more discussion of food and uid regulation as an experi

et al. 2008), while cotton nestlets may lead to conjunctivitis (Bazille et al. 2001). Bedding can also in uence mucosal immunity (Sanford et al. 2002) and endocytosis (Buddaraju and Van Dyke 2003).

Softwood beddings have been used, but the use of untreated softwood shavings and chips is contraindicated for some protocols because they can affect metabolism (Vesell 1967; Vesell et al. 1973, 1976). Cedar shav ings are not recommended because they emit aromatic hydrocarbons that induce hepatic microsomal enzymes and cytotoxicity (Torronen et al. 1989; Weichbrod et al. 1986, 1988) and have been reported to increase the inɛi dence of cancer (Jacobs and Dieter 1978; Vlahakis 1977). Prior treatment with high heat (kiln drying or autoclaving) may, depending on the material and the concentration of aromatic hydrocarbon constituents, reduce the concentration of volatile organic compounds, but the amounts remaining may be suf cient to affect speci c protocols (Cunliffe-Beamer et al. 1981; Nevalainen and Vartiainen 1996).

The purchase of bedding products should take into consideration iffeff3(s)nMk7(t
ci-3(fn-3( )-235a)-3(r)-235a166(t)-3(h)-3(h)-3(o)-3(o)-3(n)-3( )-onsidh((t)-)-3(n)-3(t)

create the need for sanitation; and the rate of soiling of the surfaces of the enclosure. Some housing systems or experimental protocols may require speci c husbandry techniques, such as aseptic handling or modi cation in the frequency of bedding change.

Agents designed to mask animal odors should not be used in animal housing facilities. They cannot substitute for good sanitation practices or for the provision of adequate ventilation, and they expose animals to volatile compounds that might alter basic physiologic and metabolic processes.

Bedding/Substrate Change Soiled bedding should be removed and replaced with fresh materials as often as necessary to keep the animals clean and dry and to keep pollutants, such as ammonia, at a concentration below levels irritating to mucous membranes. The frequency of bedding change depends on multiple factors, such as species, number, and size of the animals in the primary enclosure; type and size of the enclosure; macro- and microenvironmental temperature, relative humidity, and direct ventilation of the enclosure; urinary and fecal output and the appearance and wetness of bedding; and experimental conditions, such as those of surgery or debilitation, that might limit an animal's movement or access to clean bedding. There is no absolute minimal frequency of bedding changes; the choice is a matter of professional judgment and consultation between the investigator and animal care personnel. It typically varies from daily to weekly. In some instances frequent bedding changes are contraindicated; examples include portions of the pre- or postpartum period, research objec tives that will be affected, and species in which scent marking is critical and successful reproduction is pheromone dependent.

Cleaning and Disinfection of the Microenvironment The frequency of sanitation of cages, cage racks, and associated equipment (e.g., feeders and watering devices) is governed to some extent by the types of caging and husbandry practices used, including the use of regularly changed contact or noncontact bedding, regular ushing of suspended catch pans, and the use of wire-bottom or perforated-bottom cages. In general, enclosures and accessories, such as tops, should be sanitized at least once every 2 weeks.

2

2

appropriate, chemical agents to destroy microorganisms. Cleaning with ultrasound may be a useful method for small pieces of equipment.

If automated watering systems are used, some mechanism to ensure that microorganisms and debris do not build up in the watering devices is recommended (Meier et al. 2008); the mechanism can be periodic ushing with large volumes of water or appropriate chemical agents followed by a thorough rinsing. Constant recirculation loops that use properly maintained lters, ultraviolet lights, or other devices to disinfect recirculated water are also effective. Attention should be given to the routine sanitation of auto matic water delivery valves (i.e., lixits) during primary enclosure cleaning.

Conventional methods of cleaning and disinfection are adequate for most animal care equipment. However, it may be necessary to also sterilize caging and associated equipment to ensure that pathogenic or opportunistic microorganisms are not introduced into speci c-pathogen-free or immuno

sole means of assessing the effectiveness of the sanitation program. A decision to alter the frequency of cage bedding changes or cage washing should be based on such factors as ammonia concentration, bedding condition, appearance of the cage and animals, and the number and size of animals housed in the cage.

Mechanical washer function should be evaluated regularly and include examination of mechanical components such as spray arms and moving headers as well as spray nozzles to ensure that they are functioning appropriately. If sanitation is temperature dependent, the use of temperature-sensing devices (e.g., thermometers, probes, or temperature-sensitive indicator strips) is recommended to ensure that the equipment being sanitized is exposed to the desired conditions.

Whether the sanitation process is automated or manual, regular evaluation of sanitation effectiveness is recommended. This can be performed by evaluating processed materials by microbiologic culture or the use of organic material detection systems (e.g., adenosine triphosphate [ATP] bioluminescence) and/or by confirming the removal of artificial soil applied to equipment surfaces before washing.

Waste Disposal Conventional, biologic, and hazardous waste should be removed and disposed of regularly and safely (Hill 1999). There are several options for effective waste disposal. Contracts with licensed commercial waste disposal firms usually provide some assurance of regulatory compliance and safety. On-site incineration should comply with all federal, state, and local regulations (Nadelkov 1996).

Adequate numbers of properly labeled waste receptacles should be strategically placed throughout the facility. Waste containers should be leakproof and equipped with tight-fitting lids. It is good practice to use disposable liners and to wash containers and implements regularly. There should be a dedicated waste storage area that can be kept free of insects and other vermin. If cold storage is used to hold material before disposal, a properly labeled, dedicated refrigerator, freezer, or cold room should be used that is readily sanitized.

Hazardous wastes must be rendered safe by sterilization, containment, or other appropriate means before their removal from the facility (DHHS 2009 or most recent edition; NRC 1989, 1995b). Radioactive wastes should be kept in properly labeled containers and their disposal closely coordinated with radiation safety specialists in accord with federal and state regulations; the federal government and most states and municipalities have regulations controlling disposal of hazardous wastes. Compliance with regulations concerning hazardous-agent use (see Chapter 2) and disposal is an institutional responsibility.

Infectious animal carcasses can be incinerated on site or collected by a licensed contractor. Use of chemical digesters (alkaline hydrolysis treat

ment) may be considered in some situations (Kaye et al. 1998; Murphy et al. 2009). Procedures for on-site packaging, labeling, transportation, and storage of these wastes should be integrated into occupational health and safety policies (Richmond et al. 2003).

Hazardous wastes that are toxic, carcinogenic, flammable, corrosive, reactive, or otherwise unstable should be placed in properly labeled containers and disposed of as recommended by occupational health and safety specialists. In some circumstances, these wastes can be consolidated or blended. Sharps and glass should be disposed of in a manner that will prevent injury to waste handlers.

Pest Control Programs designed to prevent, control, or eliminate the presence of or infestation by pests are essential in an animal environment. A regularly scheduled and documented program of control and monitoring should be implemented. The ideal program prevents the entry of vermin and eliminates their harborage in the facility (Anadon et al. 2009; Easterbrook et al. 2008). For animals in outdoor facilities, consideration should be given to eliminating or minimizing the potential risk associated with pests and predators.

Pesticides can induce toxic effects on research animals and interfere with experimental procedures (Gunasekara et al. 2008). They should be used in animal areas only when necessary and investigators whose animals may be exposed to them should be consulted beforehand. Use of pesticides should be recorded and coordinated with the animal care management staff and be in compliance with federal, state, or local regulations. Whenever possible, nontoxic means of pest control, such as insect growth regulators (Donahue et al. 1989; Garg and Donahue 1989; King and Bennett 1989; Verma 2002) and nontoxic substances (e.g., amorphous silica gel), should be used. If traps are used, methods should be humane; traps that catch pests alive require frequent observation and humane euthanasia after capture (Mason and Littin 2003; Meerburg et al. 2008).

Emergency, Weekend, and Holiday CareAnimals should be cared for by qualified personnel every day, including weekends and holidays, both to safeguard their well-being and to satisfy research requirements. Emergency veterinary care must be available after work hours, on weekends, and on holidays.

In the event of an emergency, institutional security personnel and fire or police officials should be able to reach people responsible for the animals. Notification can be enhanced by prominently posting emergency procedures, names, or telephone numbers in animal facilities or by placing them in the security department or telephone center. Emergency procedures for handling special facilities or operations should be prominently posted and personnel trained in emergency procedures for these areas. A disaster plan

that takes into account both personnel and animals should be prepared as part of the overall safety plan for the animal facility. The colony manager or veterinarian responsible for the animals should be a member of the appropriate safety committee at the institution, an "official responder" in the institution, and a participant in the response to a disaster (Vogelweid 1998).

Population Management

Identification Animal records are useful and variable, ranging from limited information on identification cards to detailed computerized records for individual animals (Field et al. 2007). Means of animal identification include room, rack, pen, stall, and cage cards with written, bar-coded, or radio frequency identification (RFID) information. Identification cards should include the source of the animal, the strain or stock, names and contact information for the responsible investigator(s), pertinent dates (e.g., arrival date, birth date, etc.), and protocol number when applicable. Genotype information, when applicable, should also be included, and consistent, unambiguous abbreviations should be used when the full genotype nomenclature (see below) is too lengthy.

In addition, the animals may wear collars, bands, plates, or tabs or be marked by colored stains, ear notches/punches and tags, tattoos, subcutaneous transponders, and freeze brands. As a method of identification of small rodents, toe-clipping should be used only when no other individual identification method is feasible. It may be the preferred method for neonatal mice up to 7 days of age as it appears to have few adverse effects on behavior and well-being at this age (Castelhano-Carlos et al. 2010; Schaefer et al. 2010), especially if toe clipping and genotyping can be combined. Under all circumstances aseptic practices should be followed. Use of anesthesia or analgesia should be commensurate with the age of the animals (Hankenson et al. 2008).

Recordkeeping Records containing basic descriptive information are essential for management of colonies of large long-lived animals and should be maintained for each animal (Dyke 1993; Field et al. 2007; NRC 1979a). These records often include species, animal identifier, sire and/or dam identifier, sex, birth or acquisition date, source, exit date, and final disposition. Such animal records are essential for genetic management and historical assessments of colonies. Records of rearing and housing histories, mating histories, and behavioral profiles are useful for the management of many species, especially nonhuman primates (NRC 1979a). Relevant recorded information should be provided when animals are transferred between institutions.

Medical records for individual animals can also be valuable, especially for dogs, cats, nonhuman primates, and agricultural animals (Suckow and Doerning 2007). They should include pertinent clinical and diagnostic information,

date of inoculations, history of surgical procedures and postoperative care, information on experimental use, and necropsy findings where applicable.

Basic demographic information and clinical histories enhance the value of individual animals for both breeding and research and should be readily accessible to investigators, veterinary staff, and animal care staff.

Breeding, Genetics, and Nomenclature Genetic characteristics are important with regard to the selection and management of animals for use in breeding colonies and in biomedical research (see Appendix A). Pedigree information allows appropriate selection of breeding pairs and of experi-

temperature. The physical environment of the secondary enclosure, such as a room, constitutes themacroenvironment.

Water Quality

The composition of the water (water quality) is essential to aquatic animal well-being, although other factors that affect terrestrial microenvironments are also relevant. Water quality parameters and life support systems for aquatic animals will vary with the species, life stage, the total biomass supported, and the animals' intended use (Blaustein et al. 1999; Fisher 2000; Gresens 2004; Overstreet et al. 2000; Schultz and Dawson 2003). The success and adequacy of the system depend on its ability to match the laboratory habitat to the natural history of the species (Godfrey and Sanders 2004; Green 2002; Lawrence 2007; Spence et al. 2008).

Characteristics of the water that may affect its appropriateness include temperature, pH, alkalinity, nitrogen waste products (ammonia, nitrite, and nitrate), phosphorus, chlorine/bromine, oxidation-reduction potential, conductivity/salinity, hardness (osmolality/dissolved minerals), dissolved oxygen, total gas pressure, ion and metal content, and the established microbial ecology of the tank. Water quality parameters can directly affect animal well-being; different classes, species, and ages in a species may have different water quality needs and sensitivities to changes in water quality parameters.

Routine meT(e)-36-3(u)-3(r)-e(a)-3(m)-3(e)-(e)-3(c)-3(t)--4( )-226(a)-4(n)-32(t)-26
p1.2 Td [(t)-8(e)-8(s)-3( )-104ne s, (q)-74sti(q)-7(s)-3(s)-3( )-40()-3(t)-3(i)-3(a)-3(l)-

or neutralized before use in aquatic systems (Tompkins and Tsai 1976; Wedemeyer 2000).

Life Support System

The phrase*life support system*refers to the physical structure used to contain the water and the animals as well as the ancillary equipment used to move and/or treat the water. Life support systems may be simple (e.g., a container to hold the animal and water) or extremely complex (e.g., a fully automated recirculating system). The type of life support system used depends on several factors including the natural habitat of the species, age/size of the species, number of animals maintained, availability and characteristics of the water required, and the type of research.

Life support systems typically fall into three general categories: re‹ir culating systems where water (all or part) is moved around a system, ow-through systems where water is constantly replaced, or static systems where water is stationary and periodically replenished or replaced. The water may be fresh, brackish, or salt and is maintained at speci c temperatures depending on the species' needs.

The source of water for these systems typically falls into four general categories: treated wastewater (e.g., municipal tap water), surface water (e.g., rivers, lakes, or oceans), protected water (e.g., well or aquifer water), or arti cial water (e.g., reverse osmosis or distilled water). Arti cial saltwater may be created by adding appropriate salt to freshwater sources. Source water selection should be based on the provision of a consistent or constant supply, incoming biosecurity level requirements, water volumes needed, species selection, and research considerations.

Recirculating systems are common in indoor research settings where high-density housing systems are often needed. Most recirculating systems are designed to exchange a speci c volume of water per unit time and pe‹ri odically introduce fresh water into the system. These systems are the most mechanically advanced, containing biologic lters (bio lters) that promote conversion of ammonia to nitrite and nitrate via nitrifying bacteria, protein skimmers (foam fractionators) and particulate lters to remove undissolved and dissolved proteins and particulate matter, carbon lters to remove dissolved chemicals, and ultraviolet or ozone units to disinfect the water. The systems generally contain components to aerate and degas the water (to prevent gas oversaturation) and to heat or cool it, as well as automated dosing systems to maintain appropriate pH and conductivity. Not all ele ments are present in all systems and some components may accomplish multiple functions. Recirculating systems may be designed so that multiple individual tanks are supplied with treated water from a single source, as is the case with "rack" systems used for zebra sh (*Danio rerio*) and *Xenopus*

pensate for these thermal and moisture loads. Macroenvironmental relative humidity levels are generally de ned by safety issues and staff comfort,

these may cause subclinical effects (Smith et al. 2007). Vibration through oors can be reduced by using isolation pads under aquaria racks. Some facilities elect to place major components of the life support system (e.g., lters, pumps, and bio lters) outside the animal rooms to reduce vibration and noise.

Aquatic Housing

Microenvironment (Primary Enclosure)

The primary enclosure (a tank, raceway, pond, or pen holding water and the animal) de nes the limits of an animal's immediate environment. In research settings, acceptable primary enclosures

- allow for the normal physiological and behavioral needs of the animals, including excretory function, control and maintenance of body temperature, normal movement and postural adjustments, and, where indicated, reproduction. In some poikilothermic reptiles and amphibians, microenvironmental temperature gradients may be needed for certain physiologic functions such as feeding and digestion.
- allow conspeci c social interactions (e.g., schooling in sh species).
- provide a balanced, stable environment that supports the animal's physiologic needs.
- provide the appropriate water quality and characteristics, and per mit monitoring, lling, re lling, and changing of water.
- allow access to adequate food and removal of food waste.
- restrict escape or accidental entrapment of animals or their appendages.
- are free of sharp edges and/or projections that could cause injury.

housing for many species. When used, enrichment should elicit species-appropriate behaviors and be evaluated for safety and utility.

Generally, schooling fish species are housed with conspecifics, and many amphibians, especially anuran species, may be group housed. Aggression in aquatic animals does occur (van de Nieuwegiessen et al. 2008; Speedie and Gerlai 2008) and, as for terrestrial animals, appropriate monitoring and intervention may be necessary (Matthews et al. 2002; Torreilles and Green 2007). Some species need appropriate substrate (e.g., gravel) to reproduce or need substrate variety to express basic behaviors and maintain health (Overstreet et al. 2000). Improved breeding success in enriched environments has been reported but further research in this area is needed (Carfagnini et al. 2009). For many species (including, e.g., *X. laevis*), visual barriers, hides, and shading are appropriate (Alworth and Vasquez 2009; Torreilles and Green 2007). Most semiaquatic reptiles spend some time on land (basking, feeding, digesting, and ovipositing) and terrestrial areas should be provided as appropriate.

Sheltered, Outdoor, and Naturalistic Housing

Animals used in aquaculture are often housed in situations that mimic agricultural rearing and may be in outdoor and/or sheltered raceways, ponds, or pens with high population densities. In these settings, where natural predation and mortalities occur, it may be appropriate to measure animal "numbers" by using standard aquaculture techniques such as final production biomass (Borski and Hodson 2003).

Space

Space recommendations and housing density vary extensively with the species, age/size of the animals, life support system, and type of research (Browne et al. 2003; Green 2009; Gresens 2004; Hilken et al. 1995; Matthews et al. 2002). In the United States, for example, adult zebrafish (*Danio rerio*) in typical biomedical research settings are generally housed 5 adult fish per liter of water (Matthews et al. 2002), but this housing density varies when breeding and for housing younger animals (Matthews et al. 2002). This guidance is not necessarily relevant for other species of fish, and may change as research advances (Lawrence 2007). *X. laevis* adults may be housed at 2 liters of water per frog (NRC 1974), but a wide variety of housing systems are currently used in research settings (Green 2009). Institutions, investigators, and IACUC members should evaluate the appropriate needs of each species during program evaluations and facility inspections and continue to review ongoing research in these areas.

Aquatic Management

Behavior and Social Management

Visual evaluations of aquatic and semiaquatic animals are typically used for monitoring. To avoid damage to the protective mucus layers of the skin and negative effects on immune function (De Veer et al. 2007; Subramanian et al. 2007; Tsutsui et al. 2005), handling of these species should be kept to the minimum required (Bly et al. 1997). Appropriate handling techniques vary widely depending on the species, age/size, holding system, and specific research need (Fisher 2000; Matthews et al. 2002; Overstreet et al. 2000); they should be identified at the facility or individual protocol level.

Latex gloves have been associated with toxicity in some amphibians (Gutleb et al. 2001). The use of appropriate nets by well-trained personnel can reduce skin damage and thus stress. Nets should be cleaned and disinfected appropriately when used in different systems and should be dedicated to animals of similar health status whenever possible.

Exercise and activity levels for aquatic species are minimally described but informed decisions may be extrapolated from studies of behavior of the same or similar species in the wild (Spence et al. 2008). Some aquatic species do not rest and constantly swim; others may rest all or a significant portion of the day. Water flow rates and the provision of hides or terrestrial resting platforms (e.g., for some reptiles and amphibians) need to be appropriate for species and life stage.

Husbandry

Food The general principles relating to feeding of terrestrial animals are applicable to aquatic animals. Food should be stored in a type-appropriate manner to preserve nutritional content, minimize contamination, and prevent entry of pests. Food delivery methods should ensure that all animals are able to access food for a sufficient period of time while minimizing feeding aggression and nutrient loss. Feeding methods and frequency vary widely depending on the species, age/size of species, and type of life support system. Many aquatic or semiaquatic species are not provided with food ad libitum in the tank, and in some cases may not be fed daily.

Commercial diets (e.g., pellets, flakes) are available for certain species and storage time should be based on manufacturer recommendations or follow commonly accepted practices. In aquatic systems, particularly in fish rearing or when maintaining some amphibian and reptile species, the use of live foods (e.g., *Artemia* sp. larva, crickets, or mealworm beetle larvae) is common. Live food sources need to be maintained and managed to ensure a

steady supply and the health and suitability of the organism as a food. Care should be taken to feed a complete diet to avoid nutritional de ciencies.

Water (see also section on Water Quality)Aquatic animals need access to

Disinfection is usually accomplished through water treatment (e.g., ltration and application of UV light or ozone) and/or water changes. Chlorine and most chemical disinfectants are inappropriate for aquatic systems containing animals as they are toxic at low concentrations; when used to disinfect an entire system or system components, extreme care must be taken to ensure that residual chlorine, chemical, and reactive byproducts are neutralized or removed. The type of monitoring and frequency varies depending on the disinfection method, the system, and the animals.

Algal growth is common in aquatic systems and increases with the presence of nitrogen and phosphorus, particularly in the presence of light. Excessive growth may be an indication of elevated nitrogen or phosphorus levels. Algal species seen with recirculating systems are generally nontoxic, although species capable of producing toxins exist. Algae are typically removed using mechanical methods (i.e., scrubbing or scraping). Limiting algal growth is important to allow viewing of the animals in the enclosure. Cyanobacteria (commonly called blue-green algae) growth is also possible and may be common in freshwater aquaculture. The same factors that promote algae growth also promote cyanobacteria growth. As with algae, while most species are harmless, some species can produce clinically relevant toxic compounds (Smith et al. 2008).

Tank (cage) changing and disinfection are conducted at frequencies using methods that often differ from terrestrial systems. Because waste is dissolved in the water and/or removed as solids by siphoning or ltration, regular changing of tanks is not integral to maintaining adequate hygiene in typical aquatic systems. The frequency of cleaning and disinfection should be determined by water quality, which should permit adequate viewing of the animals, and animal health monitoring. System components such as lids on sh tanks, which may accumulate feed, may require sanitation as often as weekly depending on the frequency and type of feed and the system's design.

Cleaning and Disinfection of the Macroenvironment As with terrestrial systems, all components of the animal facility, including animal rooms and support spaces (e.g., storage areas, cage-washing facilities, corridors, and procedure rooms), should be regularly cleaned and disinfected as appropriate to the circumstances and at a frequency determined by the use of the area and the nature of likely contamination. Cleaning agents should be chosen and used with care to ensure there is no secondary contamination of the aquatic systems.

Cleaning implements should be made of materials that resist corrosion and withstand regular sanitation. They should be assigned to speci c areas and should not be transported between areas with different risks of contamination without prior disinfection. Worn items should be replaced regularly.

Waste Disposal Wastewater treatment and disposal may be necessary in some facilities depending on water volume, quality, and chemical constituents. Local regulations may limit or control the release of wastewater.

Pest Control Terrestrial animal pest control principles apply to aquatic systems but, due to transcutaneous absorption, aquatic and semiaquatic species may be more sensitive to commonly used pest control agents than terrestrial animals. Before use, an appropriate review of chemicals and methods of application is necessary.

Emergency, Weekend, and Holiday Care As with terrestrial species, aquatic animals should receive daily care from quali ed personnel who have a sufficient understanding of the housing system to identify malfunctions and, if they are unable to address a system failure of such magnitude that it requires resolution before the next workday, access to staff who can respond to the problem. Automated monitoring systems are available and may be appropriate depending on system size and complexity. Appropriate emergency response plans should be developed to address major system failures.

Population Management

Identi cation Identi cation principles are similar to those for terrestrial animals. Identi cation criteria are based on the species and housing system. Identi cation methods available for use in aquatic species include n clipping, genetic testing (Matthews et al. 2002; Nickum et al. 2004), identi ca-

illnesses; animal transfers within and out of the facility; and fertilization/hatching information (Koerber and Kalishman 2009; Matthews et al. 2002). Records should be kept concerning feeding information (e.g., food offered, acceptance), nonexpired food supplies to ensure sustenance of nutritional profile, and any live cultures (e.g., hatch rates and information to ensure suppliers' recommendations are being met; Matthews et al. 2002).

Records of water quality testing for system and source water and maintenance activities of the life support system components are important for tracking and ensuring water quality. The exact water quality parameters tested and testing frequency should be clearly established and will vary with such factors as the type of life support system, animals, and research, as discussed under Water Quality. Detailed tracking of animal numbers in aquatic systems is often possible with accurate records of transfers, breeding, and mortalities (Matthews et al. 2002). In some cases where animals are housed in large groups (e.g., some *Xenopus* colonies) periodic censuses may be undertaken to obtain an exact count. In large-scale aquaculture research it may be more appropriate to measure biomass of the system versus actual numbers of animals (Borski and Hodson 2003).

Baldwin AL. 2007. Effects of noise on rodent physiology. Int J Comp Psychol 20:134-144.
Barnett SA. 1965. Adaptation of mice to cold. Biol Rev 40:5-51.
Barnett SA. 1973. Maternal processes in the cold-adaptation of mice. Biol Rev 48:477-508.
Bartolomucci A, Palanza P, Parmigiani S. 2002. Group housed mice: Are they really stressed? Ethol Ecol Evol 14:341-350.
Bartolomucci A, Palanza P, Sacerdote P, Ceresini G, Chirieleison A, Panera AE, Parmigiani S. 2003. Individual housing induces altered immuno-endocrine responses to psychological stress in male mice. Psychoneuroendocrinology 28:540-558.
Baumans V. 1997. Environmental enrichment: Practical applications. In: Van Zutphen LFM, Balls M, eds. Animal Alternatives, Welfare and Ethics. Elsevier. p 187-197.
Baumans V. 2005. Environmental enrichment for laboratory rodents and rabbits: Requirements of rodents, rabbits, and research. ILAR J 46:162-170.
Baumans V, Schlingmann F, Vonck M, Van Lith HA. 2002. Individually ventilated cages: Be ne cial for mice and men? Contemp Top Lab Anim Sci 41:13-19.
Bayne KA. 2002. Development of the human-research animal bond and its impact on animal well-being. ILAR J 43:4-9.
Bayne KA. 2005. Potential for unintended consequences of environmental enrichment for laboratory animals and research results. ILAR J 46:129-139.
Bayne KA, Dexter SL, Hurst JK, Strange GM, Hill EE. 1993. Kong Toys for laboratory primates: Are they really an enrichment or just fomites? Lab Anim Sci 43(1):78-85.
Bayne KA, Haines MC, Dexter SL, Woodman D, Evans C. 1995. Nonhuman primate wounding prevalence: A retrospective analysis. Lab Anim 24:40-43.
Bazille PG, Walden SD, Koniar BL, Gunther R. 2001. Commercial cotton nesting material as a predisposing factor for conjunctivitis in athymic nude mice. Lab Anim (NY) 30:40-42.
Beaumont S. 2002. Ocular disorders of pet mice and rats. Vet Clin North Am Exot Anim Pract 5:311-324.
Becker BA, Christenson RK, Ford JJ, Nienaber JA, DeShazer JA, Hahn GL. 1989. Adrenal and behavioral responses of swine restricted to varying degrees of mobility. Physiol Behav 45:1171-1176.
Bell GC. 2008. Optimizing laboratory ventilation rates. Labs for the 21st century: Best practice guide. US Environmental Protection Agency. Available at http://labs21century.gov/pdf/bp_opt_vent_508.pdf; accessed March 30, 2010.
Bellhorn RW. 1980. Lighting in the animal environment. Lab Anim Sci 30:440-450.
Bergmann P, Militzer K, Büttner D. 1994. Environmental enrichment and aggressive beha viour: in uence on body weight and body fat in male inbred HLG mice. J Exp Anim Sci 37:59-78.
Berson DM, Dunn FA, Takao M. 2002. Phototransduction by retinal ganglion cells that set the circadian clock. Science 295:1070-1073.
Besch EL. 1980. Environmental quality within animal facilities. Lab Anim Sci 30:385-406.
Blaustein A, Marco A, Quichano C. 1999. Sensitivity to nitrate and nitrite in pond-breeding amphibians from the Paci c Northwest, USA. Environ Toxicol Chem J 18:2836-2839.
Blom HJM, Van Tintelen G, Van Vorstenbosch CJ, Baumans V, Beynen AC. 1996. Preferences of mice and rats for types of bedding material. Lab Anim 30:234-244.
Bloomsmith MA, Stone AM, Laule GE. 1998. Positive reinforcement training to enhance the voluntary movement of group-housed chimpanzees within their enclosures. Zoo Biol 17:333-341.
Bly JE, Quiniou SM, Clem LW. 1997. Environmental effects on sh immune mechanisms. Dev Biol Stand 90:33-43.
Borski R, Hodson RG. 2003. Fish research and the institutional animal care and use committee. ILAR J 44:286-294.

Bracke MBM, Metz JHM, Spruijt BM, Schouten WGP. 2002. Decision support system for overall welfare assessment in pregnant sows. B: Validation by expert opinion. J Anim Sci 80:1835-1845.

Brainard GC. 1989. Illumination of laboratory animal quarters: Participation of light irradiance and wavelength in the regulation of the neuroendocrine system. In: Science and Animals: Addressing Contemporary Issues. Greenbelt, MD: Scientists Center for Animal Welfare. p 69-74.

Brainard GC, Vaughan MK, Reiter RJ. 1986. Effect of light irradiance and wavelength on the Syrian hamster reproductive system. Endocrinology 119:648-654.

Brenner FJ, Brenner PE. 1969. The in uence of light and temperature on body fat and repro ductive conditions of Rana pipiens. Ohio J Sci 69:305-MC /ref <<uc

)-3(8)-3(6)43(.)-3( )-80(9)-3(f)-3(f)-3(e)-3(d)-3(t)-3( )-65(9)-3(f)-3( )-52(9)p3(a)-3(r)-3(e)-3( )-52(9)-3(.)-3(d)u3(n)-3(d)-3( )-52

Br9T-319TL31r9T-319TC(r)-3o9T-319T13o9TTr9T-3or9To3o9Te31r9Ta319Ti319T-3o

Bred s A3(.)M(J)S-3( )-52116ee s

Shaston 0<B3(t)-3(o)-3(n)c3(h)-3(i)-3(m)-3(p)-31(S)6c3-MC /r73(9)3f <<3( )]TJ /T1_0

SreFJc-31(S)-3(n)-3(d)-3( )-52 1rn FJc-31(S)b3(e)-3(l)-3(e)l3( )-31(O)f3(r)-3(e)-3

S3c -M4B3c -M4B

Bred ss aJ edFJeaah J atFJe5

Snd are tat Of Rtoireat0ure 5.763 Td [(R)t3( )aT( )d3(a)p3(a)-3(n)l3( )e3(d)

Chmiel DJ, Noonan M. 1996. Preference of laboratory rats for potentially enriching stimulus objects. Lab Anim 30:97-101.

Clarence WM, Scott JP, Dorris MC, Paré M. 2006. Use of enclosures with functional vertical space by captive rhesus monkeys (Macaca mulatta) involved in biomedical research. JAALAS 45:31-34.

Georgsson L, Barrett J, Gietzen D. 2001. The effects of group-housing and relative weight on feeding behaviour in rats. Scand J Lab Anim Sci 28:201-209.
Gibson SV, Besch-Williford C, Raisbeck MF, Wagner JE, McLaughlin RM. 1987. Organophosphate toxicity in rats associated with contaminated bedding. Lab Anim 37:789-791.
Gill TJ. 1980. The use of randomly bred and genetically de ned animals in biomedical research. Am J Pathol 101(3S):S21-S32.
Godfrey EW, Sanders GW. 2004. Effect of water hardness on oocyte quality and embryo development in the African clawed frog (Xenopus laevis). Comp Med 54:170-175.
Gonder JC, Laber K. 2007. A renewed look at laboratory rodent housing and management. ILAR J 48:29-36.
Gonzalez RR, Kiuger MJ, Hardy JD. 1971. Partitional calorimetry of the New Zealand white rabbit at temperatures of 5-35°C. J Appl Physiol 31:728.
Gordon AH, Hart PD, Young MR. 1980. Ammonia inhibits phagosome-lysosome fusion in macrophages. Nature 286:79-80.
Gordon CJ. 1990. Thermal biology of the laboratory rat. Physiol Behav 47:963-991.
Gordon CJ. 1993. Temperature Regulation in Laboratory Animals. New York: Cambridge University Press.
Gordon CJ. 2004. Effect of cage bedding on temperature regulation and metabolism of group-housed female mice. Comp Med 54:63-68.
Gordon CJ. 2005. Temperature and Toxicology: An integrative, comparative and environmental approach. Boca Raton, FL: CRC Press.
Gordon CJ, Becker P, Ali JS. 1998. Behavioral thermoregulatory responses of single- and group-housed mice. Physiol Behav 65:255-262.
Green EL. 1981. Genetics and Probability in Animal Breeding Experiments. New York: Oxford University Press.
Green SL. 2002. Factors affecting oogenesis in the South African clawed frog (Xenopus laevis). Comp Med 52:307-312.
Green SL. 2009. The Laboratory Xenopus sp. (Laboratory Animal Pocket Reference). Boca Raton, FL: CRC Press.
Greenman DL, Bryant P, Kodell RL, Sheldon W. 1982. In uence of cage shelf level on retinal atrophy in mice. Lab Anim Sci 32:353-356.
Gresens J. 2004. An introduction to the Mexican axolotl (Ambystoma mexicanum). Lab Anim 33:41-47.
Groen A. 1977. Identi cation and genetic monitoring of mouse inbred strains using biomedical polymorphisms. Lab Anim (London) II:209-214.
Gunasekara AS, Rubin AL, Goh KS, Spurlock FC, Tjeerdema RS. 2008. Environmental fate and toxicology of carbaryl. Rev Environ Contam Toxicol 196:95-121.
Gutleb AC, Bronkhorst M, van den Berg JHJ, Musrk AJ. 2001. Latex laboratory gloves: An unexpected pitfall in amphibian toxicity assays with tadpoles. Environ Toxicol Pharmacol 10:119-121.
Haemisch A, Voss T, Gärtner K. 1994 Effects of environmental enrichment on aggressive behaviour, dominance hierarchies and endocrine states in male DBA/2J mice. Physiol Behav 56:1041-1048.
Hahn NE, Lau D, Eckert K, Markowitz H. 2000. Environmental enrichment-related injury in a macaque (Macaca fascicularis): Intestinal linear foreign body. Comp Med 50:556-558.
Hall FS. 1998. Social deprivation of neonatal, adolescent, and adult rats has distinct neurochemical and behavioural consequences. Crit Rev Neurobiol 12:129-162.
Hall JE, White WJ, Lang CM. 1980. Acidi cation of drinking water: Its effects on selected biologic phenomena in male mice. Lab Anim Sci 30:643-651.

Hankenson FC, Garzel LM, Fischer DD, Nolan B, Hankenson KD. 2008. Evaluation of tail biopsy collection in laboratory mice (Mus musculus): Vertebral ossification, DNA quantity, and acute behavioral responses. JAALAS 47(6):10-18.

Hanifin JP, Brainard GC. 2007. Photoreception for circadian, neuroendocrine, and neurobehavioral regulation. J Physiol Anthropol 26:87-94.

Hartl DL. 2000. A Primer of Population Genetics, 3rd ed. Sunderland, MA: Sinauer Associates.

Hasenau JJ, Baggs RB, Kraus AL. 1993. Microenvironments in microisolation cages using BALB/c and CD-1 Mice. Contemp Top Lab Anim Sci 32:11-16.

Hedrich HJ. 1990. Genetic Monitoring of Inbred Strains of Rats. New York: Gustav Fischer Verlag.

Heffner HE, Heffner RS. 2007. Hearing ranges of laboratory animals. JAALAS 46:20-22.

Held SDE, Turner RJ, Wootton RJ. 1995. Choices of laboratory rabbits for individual or group-housing. Appl Anim Behav Sci 46:81-91

Hermann LM, White WJ, Lang CM. 1982. Prolonged exposure to acid, chlorine, or tetracycline in drinking water: Effects on delayed-type hypersensitivity, hemagglutination titers, and reticuloendothelial clearance rates in mice. Lab Anim Sci 32:603-608.

Hess SE, Rohr S, Dufour BD, Gaskill BN, Pajor EA, Garner JP. 2008. Home improvement: C57BL/6J mice given more naturalistic nesting materials build better nests. JAALAS 47:25-31.

Hilken G, Dimigen J, Iglauer F. 1995. Growth of Xenopus laevis under different laboratory rearing conditions. Lab Anim 29:152-162.

Hill D. 1999. Safe handling and disposal of laboratory animal waste. Occup Med 14:449-468.

Hoffman HA, Smith KT, Crowell JS, Nomura T, Tomita T. 1980. Genetic quality control of laboratory animals with emphasis on genetic monitoring. In: Spiegel A, Erichsen S, Solleveld HA, eds. Animal Quality and Models in Biomedical Research. Stuttgart: Gustav Fischer Verlag. p 307-317.

Homberger FR, Pataki Z, Thomann PE. 1993. Control of Pseudomonas aeruginosa infection in mice by chlorine treatment of drinking water. Lab Anim Sci 43:635-637.

Hotchkiss CE, Paule MG. 2003. Effect of pair-housing on operant behavior task performance

Kaye GI, Weber PB, Evans A, Venezia RA. 1998. Ef cacy of alkaline hydrolysis as an alterna tive method for treatment and disposal of infectious animal waste. Contemp Top Lab Anim Sci 37:43-46.

Keenan KP, Smith PF, Soper KA. 1994. Effect of dietary (caloric) restriction on aging, survival, pathobiology and toxicology. In: Notter W, Dungworth DL, Capen CC, eds. Pathobiology of the Aging Rat, vol 2. Washington: International Life Sciences Institute. p 609-628.

Keenan KP, Laroque P, Ballam GC, Soper KA, Dixit R, Mattson BA, Adams SP, Coleman JB. 1996. The effects of diet, ad libitum overfeeding, and moderate dietary restriction on the rodent bioassay: The uncontrolled variable in safety assessment. Toxicol Pathol 24:757-768.

Keller LSF, White WJ, Snyder MT, Lang CM. 1989. An evaluation of intracage ventilation in three animal caging systems. Lab Anim Sci 39:237-242.

Kempthorne O. 1957. An Introduction to Genetic Statistics. New York: John Wiley and Sons.

King JE, Bennett GW. 1989. Comparative activity of fenoxycarb and hydroprene in sterilizing the German cockroach (Dictyoptera: Blattellidae). J Econ Entomol 82:833-838.

Liu L, Nutter LMJ, Law N, McKerlie C. 2009. Sperm freezing and in vitro fertilization on three substrains of C57BL/6 mice. JAALAS 48:939-943.

Lupo C, Fontani G, Girolami L, Lodi L, Muscettola M. 2000. Immune and endocrine aspects of physical and social environmental variations in groups of male rabbits in seminatural conditions. Ethol Ecol Evol 12:281-289.

Lutz CK, Novak MA. 2005. Environmental enrichment for nonhuman primates: Theory and application. ILAR J 46:178-191.

MacCluer JW, VandeBerg JL, Read B, Ryder OA. 1986. Pedigree analysis by computer simulation. Zoo Biol 5:147-160.

MacLean EL, Prior RS, Platt ML, Brannon EM. 2009. Primate location preference in a double-

Murphy RGL, Scanga JA, Powers BE, Pilon JL, VerCauteren KC, Nash PB, Smith GC, Belk KE. 2009. Alkaline hydrolysis of mouse-adapted scrapie for inactivation and disposal of prion-positive material. J Anim Sci 87:1787-1793.

Nadelkov M. 1996. EPA impact on pathological incineration: What will it take to comply? Lab Anim 25:35-38.

NAFA [National Air Filtration Association]. 1996. NAFA Guide to Air Filtration, 2nd ed. Virginia Beach.

NASA [National Aeronautics and Space Administration]. 1988. Summary of conclusions reached in workshop and recommendations for lighting animal housing modules used in microgravity related projects. In: Holley DC, Winget CM, Leon HA, eds. Lighting Require ments in Microgravity: Rodents and Nonhuman Primates. NASA Technical Memorandum 101077. Moffett Field, CA: Ames Research Center. p 5-8.

Nay eld KC, Besch EL. 1981. Comparative responses of rabbits and rats to elevated noise. Lab Anim Sci 31:386-390.

Nevalainen T, Vartiainen T. 1996. Volatile organic compounds in commonly used beddings before and after autoclaving. Scand J Lab Anim Sci 23:101-104.

Newberne PM. 1975. In uence on pharmacological experiments of chemicals and other fac tors in diets of laboratory animals. Fed Proc 34:209-218.

Newberry RC. 1995. Environmental enrichment: Increasing the biological relevance of captive environments. Appl Anim Beh Sci 44:229-243.

Newbold JA, Chapin LT, Zinn SA, Tucker HA. 1991. Effects of photoperiod on mammary development and concentration of hormones in serum of pregnant dairy heifers. J Dairy Sci 74:100-108.

Nickum JG, Bart HL Jr, Bowser PR. 2004. Guidelines for the Use of Fishes in Research. Bethesda, MD: American Fisheries Society.

Njaa LR, Utne F, Braek( )-40(F)-2(:)-2( )2 <</MCID 6243 >>BDCwf ll-40(L)-3(R)-3(m)-3(e)-3(e)-3(e)-3(eF)-
pRebeC EMC EM(o)-3:6243 >>38-004. Guideine fo eE.on foeboa-3((t)-3a-3(g-2(f)-3(o186(o(v)-3(e)-3

NRC. 1998a. Psychological Well-being of Nonhuman Primates. Washington: National Academy Press.
NRC. 1998b. Nutrient Requirements of Swine, 10th rev ed. Washington: National Academy Press.
NRC. 2000. Nutrient Requirements of Beef Cattle, 7th rev ed: Update 2000. Washington: National Academy Press.
NRC. 2001. Nutrient Requirements of Dairy Cattle, 7th rev ed. Washington: National Academy Press.
NRC. 2003a. Nutrient Requirements of Nonhuman Primates, 2nd rev ed. Washington: National Academies Press.
NRC. 2003b. Guidelines for the Care and Use of Mammals in Neuroscience and Behavioral Research. Washington: National Academies Press.
NRC. 2006a. Preparation of Animals for Use in the Laboratory. ILAR J 43:281-375.
NRC. 2006b. Nutrient Requirements of Dogs and Cats. Washington: National Academies Press.
NRC. 2006c. Nutrient Requirements of Horses, 6th rev ed. Washington: National Academies Press.
NRC. 2007. Nutrient Requirements of Small Ruminants: Sheep, Goats, Cervids, and New World Camelids. Washington: National Academies Press.
Olivier B, Molewijk E, van Oorschot R, van der Poel G, Zethof T, van der Heyden J, Mos J. 1994. New animal models of anxiety. Eur Neuropsychopharmacol 4:93-102.
Olson LC, Palotay JL. 1983. Epistaxis and bullae in cynomolgus macaques (Macaca fascicularis). Lab Anim Sci 33:377-379.
Olsson IA, Dahlborn, K. 2002. Improving housing conditions for laboratory mice: A review of "environmental enrichment." Lab Anim 36:243-270.
OSHA [Occupational Safety and Health Administration]. 1998. Occupational Safety and Health Standards. Subpart G, Occupational Health and Environmental Controls, Occupational Noise Exposure (29 CFR 1910.95). Washington: Department of Labor.
O'Steen WK. 1980. Hormonal influences in retinal photodamage. In: Williams TP, Baker BN, eds. The Effects of Constant Light on Visual Processes. New York: Plenum Press. p 29-49.

Perkins SE, Lipman NS. 1995. Characterization and quali cation of microenvironmental contaminants in isolator cages with a variety of contact bedding. Contemp Top Lab Anim Sci 34:93-98.
Peterson EA. 1980. Noise and laboratory animals. Lab Anim Sci 30:422-439.
Peterson EA, Augenstein JS, Tanis DC, Augenstein DG. 1981. Noise raises blood pressure without impairing auditory sensitivity. Science 211:1450-1452.
Pfaff J, Stecker M. 1976. Loudness levels and frequency content of noise in the animal house. Lab Anim 10:111-117.
Poiley SM. 1960. A systematic method of breeder rotation for non-inbred laboratory animal colonies. Proc Anim Care Panel 10:159-166.
Poole T. 1998. Meeting a mammal's psychological needs. In: Shepherdson DJ, Mellen JD, Hutchins M, eds. Second Nature: Environmental Enrichment for Captive Animals. Washington: Smithsonian Institute Press. p 83-94.
Pough FH. 1991. Recommendations for the care of amphibians and reptiles in academic institutions. ILAR J 33:1-16.
Pough FH. 2007. Amphibian biology and husbandry. ILAR J 48:203-213.
Raje S. 1997. Group housing for male New Zealand white rabbits. Lab Anim 26:36-38.
Ras T, van de Ven M, Patterson-Kane EG, Nelson K. 2002. Rats' preferences for corn versus wood-based bedding and nesting materials. Lab Anim 36:420-425.
Rasmussen S, Glickman G, Norinsky R, Quimby F, Tolwani RJ. 2009. Construction noise decreases reproductive ef ciency in mice. JAALAS 48:263-270.
Reeb CK, Jones R, Bearg D, Bedigan H, Myers D, Paigen B. 1998. Microenvironment in ventilated cages with differing ventilation rates, mice populations and frequency of bedding changes. JAALAS 37:70-74.
Reeb-Whitaker CK, Paigen B, Beamer WG, Bronson RT, Churchill GA, Schweitzer IB, Myers DD. 2001. The impact of reduced frequency of cage changes on the health of mice housed in ventilated cages. Lab Anim 35:58-73.
Reinhardt V. 1997. Training nonhuman primates to cooperate during handling procedures: A review. Anim Tech 48:55-73.
Reme CE, Wirz-Justice A, Terman M. 1991. The visual input stage of the mammalian circadian pacemaking system. I: Is there a clock in the mammalian eye? J Biol Rhythms 6:5-29.
Rennie AE, Buchanan-Smith HM. 2006. Re nement of the use of non-human primates in scienti c research. Part I: The in uence of humans. Anim Welf 15:203-213.
Richmond JY, Hill RH, Weyant RS, Nesby-O'Dell SL, Vinson PE. 2003. What's hot in animal biosafety? ILAR J 44:20-27.
Roberts-Thomson A, Barnes A, Filder DS, Lester RJG, Adlard RD. 2006. Aerosol dispersal of the sh pathogen Amyloodinium ocellatum. Aquaculture 257:118-123.
Rock FM, Landi MS, Hughes HC, Gagnon RC. 1997. Effects of caging type and group size on selected physiologic variables in rats. Contemp Top Lab Anim Sci 36:69-72.
Rollin BE. 1990. Ethics and research animals: Theory and practice. In: Rollin B, Kesel M, eds. The Experimental Animal in Biomedical Research, vol I: A Survey of Scienti c and Ethical Issues for Investigators. Boca Raton, FL: CRC Press. p 19-36.
Rommers J, Meijerhof R. 1996. The effect of different oor types on foot pad injuries of rabbit does. In: Proceedings of the 6th World Rabbit Science Congress 1996, Toulouse. p 431-436.
Russell RJ, Festing MFW, Deeny AA, Peters AG. 1993. DNA ngerprinting for genetic monitoring of inbred laboratory rats and mice. Lab Anim Sci 43:460-465.
Sales GD. 1991. The effect of 22 kHz calls and arti cial 38 kHz signals on activity in rats. Behav Proc 24:83-93.
Sales GD, Milligan SR, Khirnykh K. 1999. Sources of sound in the laboratory animal environment: A survey of the sounds produced by procedures and equipment. Anim Welf 8:97-115.

Thigpen JE, Lebetkin EH, Dawes ML, Clark JL, Langley CL, Amy HL, Crawford D. 1989. A standard procedure for measuring rodent bedding particle size and dust content. Lab Anim Sci 39:60-62.

Thigpen JE, Setchell KDR, Ahlmark KB, Locklear J, Spahr T, Caviness GF, Goelz MF, Haseman JK, Newbold RR, Forsythe DB. 1999. Phytoestrogen content of purified, open- and closed-formula laboratory animal diets. Lab Anim Sci 49:530-539.

Thigpen JE, Setchell KDR, Saunders HE, Haseman JK, Grant MG, Forsythe DB. 2004. Selecting the appropriate rodent diet for endocrine disruptor research and testing studies. ILAR J 45:401-416.

Tompkins JA, Tsai C. 1976. Survival time and lethal exposure time for the blacknose dace exposed to free chlorine and chloramines. Trans Am Fish Soc 105:313-321.

Torreilles SL, Green SL. 2007. Refuge cover decreases the incidence of bite wounds in laboratory South African clawed frogs (Xenopus laevis). JAALAS 46:33-36.

Torronen R, Pelkonen K, Karenlampi S. 1989. Enzyme-inducing and cytotoxic effects of wood-based materials used as bedding for laboratory animals: Comparison by a cell culture study. Life Sci 45:559-565.

Totten M. 1958. Ringtail in newborn Norway rats: A study of the effect of environmental temperature and humidity on incidence. J Hygiene 56:190-196.

Tsai PP, Stelzer HD, Hedrich HJ, Hackbarth H. 2003. Are the effects of different enrichment designs on the physiology and behaviour of DBA/2 mice consistent? Lab Anim 37:314-327.

Tsutsui S, Tasumi S, Suetake H, Kikuchi K, Suzuki Y. 2005. Demonstration of the mucosal lectins in the epithelial cells of internal and external body surf20s6)-163(m)-3 <</MCID 6494 >>B(i)--3 <<
Tsu9-3(t)-3(s)-3(u)-3(i)-3( )-11(S)w
ku(7)-33Ã£L3: <3ÌL:43Ó NÌCê4Ú=3¡S73Ú3£C3>ó1%3oisl
ku(7)-33n(z)-3i(a)-<</MC--3 <z

Vesell ES. 1967. Induction of drug-metabolizing enzymes in liver microsomes of mice and rats by softwood bedding. Science 157:1057-1058.

Vesell ES, Lang CM, White WJ, Passananti GT, Tripp SL. 1973. Hepatic drug metabolism in rats: Impairment in a dirty environment. Science 179:896-897.

Vesell ES, Lang CM, White WJ, Passananti GT, Hill RN, Clemen TL, Liu DL, Johnson WD. 1976. Environmental and genetic factors affecting response of laboratory animals to drugs. Fed Proc 35:1125-1132.

Vlahakis G. 1977. Possible carcinogenic effects of cedar shavings in bedding of C3H-AV/AB mice. J Natl Cancer Inst 58:149-150.

Vogelweid CM. 1998. Developing emergency management plans for university laboratory animal programs and facilities. Contemp Top Lab Anim Sci 37:52-56.

Waiblinger E. 2002. Comfortable quarters for gerbils in research institutions. In: Reinhardt V, Reinhardt A, eds. Comfortable Quarters for Laboratory Animals, 9th ed. Washington: Animal Welfare Institute. p 18-25.

Wardrip CL, Artwohl JE, Bennett BT. 1994. A review of the role of temperature versus time in an effective cage sanitation program. Contemp Top Lab Anim Sci 33:66-68.

Wardrip CL, Artwohl JE, Oswald J, Bennett BT. 2000. Veri cation of bacterial killing effects of cage wash time and temperature combinations using standard penicylinder methods. Contemp Top Lab Anim Sci 39:9-12.

Wax TM. 1977. Effects of age, strain, and illumination intensity on activity and self-selection of light-dark schedules in mice. J Comp Physiol Psychol 91:51-62.

Wedemeyer GA. 2000. Chlorination/dechlorination. In: Stickney RR, ed. Encyclopaedia of Aquaculture. Chichester: John Wiley and Sons. p 172-174.

Weed JL, Watson LM. 1998. Pair housing adult owl monkeys (Aotus sp.) for environmental enrichment. Am J Primatol 45:212.

Weichbrod RH, Hall JE, Simmonds RC, Cisar CF. 1985. Selecting bedding material. Lab Anim 15:25-29.

Weichbrod RH, Cisar CF, Miller JG, Simmonds RC, Alvares AP, Ueng TH. 1988. Effects of cage beddings on microsomal oxidative enzymes in rat liver. Lab Anim Sci 38:296-298.

Wooster GA, Bowser PR. 2007. The aerobiological pathway of a sh pathogen: Survival and disseminaton of Aeromonas salmonicida in aerosols and its implications in sh health management. J World Aquacul Soc 27:7-14.

Würbel H. 2001. Ideal homes? Housing effects on rodent brain and behaviour. Trends Neurosci 24:207-211.

Yanong RPE. 2003. Fish health management considerations in recirculating aquaculture sys tems, part 2: Pathogens. IFAS, University of Florida. Available at www.aces.edu/dept/ sh eries/aquaculture/documents/ shhealth2.pdf; accessed April 15, 2010.

Yildiz A, Hayirli A, Okumus Z, Kaynar O, Kisa F. 2007. Physiological pro le of juvenile rats: Effects of cage size and cage density. Lab Anim 36:28-38.

Young RJ. 2003. Environmental Enrichment for Captive Animals. UFAW Animal Welfare Series. London: Blackwell Science.

4

Veterinary Care

Veterinary care is an essential part of an animal care and use Program.

from these sources and the potential for introducing health risks to personnel and other facility animals. Breeding colonies should be established based on need and managed according to principles of animal reduction such as cryopreservation for rodent stocks or strains (Robinson et al. 2003).

Transportation of Animals

Transportation of animals is governed by a number of US regulatory agencies and international bodies. The Animal Welfare Regulations (USDA 1985) set standards for interstate and export/import transportation of regulated species; the International Air Transport Association (IATA) updates the Live Animals Regulations annually and IATA member airlines and many countries agree to comply with these regulations to ensure the safe and humane transport of animals by air (IATA 2009). The Centers for Disease Control and Prevention and USDA enforce regulations to prevent the introduction, transmission, or spread of communicable diseases and regulate the importation of any animal or animal product capable of carrying a zoonotic disease. The US Fish and Wildlife Service regulates importation/exportation of wild vertebrate and invertebrate animals and their tissues. As the national authority arm of the Convention on International Trade in Endangered Spe

receipt. Shipping should be coordinated to ensure that animals arrive during normal business hours or, if delivery occurs outside of this time, that someone is available to receive them. Defining and delegating responsibility to the appropriate persons, who are knowledgeable about the needs of the species being shipped, will help ensure effective communication and planning of animal transport (AVMA 2002).

All animals in transit within and between institutions or jurisdictions should be accompanied by appropriate documentation to minimize delays in shipping and receipt. Documentation may include health certificates, sending and receiving institutions' addresses and contacts, emergency procedures and veterinary contact information, and agency permits as needed.

For noncommercial sources of animals, in particular, it is important for the veterinarian or the veterinarian's designee to review the health status and other housing and husbandry requirements before authorizing shipment of animals. This action will ensure that effective quarantine practices are implemented for incoming animals and address any special requirements needed to ensure animal well-being (Otto and Tolwani 2002). Special considerations may be necessary for transporting animals during certain phases of their life or in certain conditions, such as pregnant, perinatal, and geriatric animals; animals with preexisting medical conditions (e.g., diabetes mellitus); and animals surgically prepared by the supplier (FASS 2010).

Although ensuring animal biosecurity during transportation is always important, it is of particular importance for immunocompromised, genetically modified, and specific pathogen-free rodents (Jacoby and Lindsey 1998). For these animals, reinforced disposable shipping containers with filter-protected ventilation openings and internal food and water sources help ensure that microbial contamination does not occur during transit. Commercial vendors are experienced in animal transport and typically use dedicated transport systems and protocols to minimize microbiologic contamination. Noncommercial or interinstitutional transfer of rodents poses a higher risk of microbial contamination since the individuals involved may lack the required knowledge and animal biosecurity capabilities to maintain the animals' health status. Risks due to in-transit microbial contamination of shipping container surfaces can be reduced by decontaminating the surfaces before placement of the containers in clean sites of animal facilities (NRC 1996, 2006). Transportation of animals in private vehicles is discouraged because of potential animal biosecurity, safety, health, and liability risks for the animals, personnel, and institution.

For aquatic species and amphibians, special considerations are required for transportation in an aqueous or sufficiently moist environment, and special attention should be given to avoiding temperature extremes for poikilotherms.

In all cases, appropriate loading and unloading facilities should be provided for the safe and secure transfer of animals at an institution. Facilities and procedures should be in place to help ensure that the environment at the site does not pose risks to animal well-being or personnel safety. During times of extreme temperatures animal transport may be detrimental to animal well-being and therefore may not be possible unless an appropriately heated or cooled means of transportation is available (Robertshaw 2004; Schrama et al. 1996).

PREVENTIVE MEDICINE

Disease prevention is an essential component of comprehensive veterinary medical care and biosecurity programs. Effective preventive medicine enhances the research value of animals by maintaining healthy animals and minimizing nonprotocol sources of variation associated with disease and inapparent infection, thus minimizing animal waste and potential effects on well-being. Preventive medicine programs consist of various combinations of policies, procedures, and equipment related to quarantine and stabilization and the separation of animals by species, source, and health status.

Animal Biosecurity

Animal biosecurity refers to all measures taken to identify, contain, prevent, and eradicate known or unknown infections that may cause clinical disease or alter physiologic and behavioral responses or otherwise make the animals unsuitable for research. Animal biosecurity practices should be applied to all species, but they are most important when housing large numbers of animals in intensive housing conditions (e.g., laboratory rodents). Limiting exposure of animals to infectious disease agents requires consideration of physical plant layout and operational practices. Separation of clean and soiled caging and equipment, and sometimes the associated staff, is often fundamental to success.

Animal biosecurity includes all measures to control known or unknown infections in laboratory animals.

A successful animal biosecurity program incorporates a number of elements: procedures that ensure that only animals of a desired defined health status enter the facility; personnel and materials, especially consumables, that do not serve as fomites; practices that reduce the likelihood of cross contamination if an infectious agent is inadvertently introduced; a comprehensive ongoing system for evaluating animals' health status, including access to all animals; and containment and eradication, if desired, of

introduced infectious agents. Related components include procedures for evaluating and selecting appropriate animal suppliers (these may include quarantine and determination of animal health status if unknown); treatment of animals or their products at entry to minimize disease risks (e.g., surface disinfection of fish eggs); a comprehensive pest control program that may include evaluation of the health status of feral animals; procedures to ensure that all biologics administered to animals are free of contamination; and procedures for intra- and interfacility animal transport (e.g., transport of animals to laboratory and other facilities outside the animal facility can present challenges to animal biosecurity) (Balaban and Hampshire 2001). Additional details pertaining to these topics are available in the sections of Chapter 2 that deal with occupational health and safety.

Quarantine and Stabilization

Quarantine is the separation of newly received animals from those already in the facility, in a way that prevents potential spread of contaminants, until the health and possibly the microbial status of the newly received animals have been determined. Transportation can be stressful and may induce recrudescence of subclinical infections harbored by an animal.

An effective quarantine program minimizes the risk of introduction of

Depending on the health status of the colony animals and consistent with the animal biosecurity program in place, rodents or other animals being moved outside an animal facility for procedures (e.g., imaging or behavioral testing) may need to be held separately from their colony of origin until their health status is evaluated.

Regardless of whether the animals are quarantined, newly received animals should be given a period for physiologic, behavioral, and nutritional acclimation before their use (Obernier and Baldwin 2006). The length of time for acclimation will depend on the type and duration of animal transportation, the species, and the intended use of the animals. For animals not typically housed in research settings, consideration should be given to providing means to assist with their acclimation (e.g., shearing sheep before they are brought indoors). The need for an acclimation period has been demonstrated in mice, rats, guinea pigs, nonhuman primates, and goats, and time for acclimation is likely important for other species as well (Capitanio et al. 2006; Conour et al. 2006; Kagira et al. 2007; Landi et al. 1982; Prasad et al. 1978; Sanhouri et al. 1989; Tuli et al. 1995).

Separation by Health Status and Species

Physical separation of animals by species is recommended to prevent interspecies disease transmission and to eliminate the potential for anxiety and physiologic and behavioral changes due to interspecies conflict (Arndt et al. 2010). Such separation is usually accomplished by housing different species in separate rooms, but in some instances it may be possible with cubicles, laminar flow units, cages that have filtered air or separate ventilation, or isolators. It may also be acceptable to house different species in the same room—for example, two species that have a similar pathogen status and are behaviorally compatible (Pritchett-Corning et al. 2009), or aquatic species, as long as nets and other animal handling devices remain separate between systems.

In some species subclinical or latent infections can cause clinical disease if transmitted to another species. A few examples may serve as a guide in determining the need for separate housing by species:

- Helicobacter bilis can infect rats and mice and may induce clinical disease in both species (Haines et al. 1998; Jacoby and Lindsey 1998; Maggio-Price et al. 2002).
- As a rule, New World (South and Central American), Old World African, and Old World Asian species of nonhuman primates should be housed in separate rooms. Simian hemorrhagic fever (Renquist 1990) and simian immunodeficiency virus (Hirsch et al. 1991; Murphey-Corb et al. 1986), for example, cause only subclinical

infections in African species but induce clinical disease in Asian species.

- Some species should be housed in separate rooms even though they are from the same geographic region. For example, squirrel monkeys (Saimiri sciureus) and tamarins (Saguinus oedipus) may be latently infected with herpesviruses (Herpesvirus saimiri and H. tamarinus, respectively2elys

and serology. If a disease or infectious agent is identi ed in a facility or colony, the choice of therapy should be made by the veterinarian in coп sultation with the investigator. If the animal is to remain in the study, the selected treatment plan should be therapeutically sound and, when pөs sible, interfere minimally with the research process.

Subclinical microbial infections (see Appendix A, Pathology, Clinical Pathology, and Parasitology) occur frequently in conventionally maintained rodents but can also occur in facilities designed and maintained for produc tion and use of pathogen-free rodents if the microbial barrier is breached. Examples of infectious agents that can be subclinical but that may induce immunologic changes or alter physiologic, pharmacologic, or toxicologic responses are noroviruses, parvoviruses, mouse he,y or als hp7D-3(l)0-2(s)-2( )26(a) -litio andc

program's requirements, which will vary by institution, species used, and the nature of the animal use. To be effective in providing clinical care, the veterinarian should be familiar with the species and various uses of animals in the institutional research, teaching, testing, or production programs and have access to medical and experimental treatment records.

Medical Management

There should be a timely and accurate method for communication of any abnormalities in or concerns about animal health, behavior, and well-being to the veterinarian or the veterinarian's designee. The responsibility for communicating these concerns rests with all those involved with animal care and use. Reports should be triaged to ensure that animals most in need receive priority attention, and the veterinarian or veterinarian's designee should perform an objective assessment of the animal(s) to determine an appropriate course of action.

Well-planned experiments with clearly delineated scientific and humane endpoints will help to ensure that a contingency plan is in place for problems that may arise during the study (see Chapter 2, Experimental and Humane Endpoints). For animals on research protocols, the veterinarian or veterinarian's designee should make every effort to discuss any problems with the principal investigator or project director to jointly determine the most appropriate course of treatment or action. Standard operating procedures (SOPs) may be developed for recurrent health conditions to expedite treatment. Recurrent or significant problems involving experimental animal health should be communicated to the IACUC, and all treatments and outcomes should be documented (USDA 1997).

Emergency Care

Procedures must be in place to provide for emergency veterinary care both during and outside of regularly scheduled hours. Such procedures must enable animal care and research staff to make timely reports of animal injury, illness, or death. A veterinarian or the veterinarian's designee must be available to expeditiously assess the animal's condition, treat the animal, investigate an unexpected death, or advise on euthanasia. In the case of a pressing health problem, if the responsible person (e.g., investigator) is not available or if the investigator and veterinary staff cannot reach consensus on treatment, the veterinarian must have the authority, delegated by senior administration (see Chapter 2, Institutional Official and Attending Veterinarian) and the IACUC, to treat the animal, remove it from the experiment, institute appropriate measures to relieve severe pain or distress, or perform euthanasia if necessary.

Recordkeeping

Medical records are a key element of the veterinary care program and are considered critical for documenting animal well-being as well as tracking animal care and use at a facility. A veterinarian should be involved in establishing, reviewing, and overseeing medical and animal use records (Field et al. 2007; Suckow and Doerning 2007). All those involved in animal care and use must comply with federal laws and regulations regarding human and veterinary drugs and treatments. Drug records and storage procedures should be reviewed during facility inspections.

SURGERY

Successful surgical outcomes require appropriate attention to presurgical planning, personnel training, anesthesia, aseptic and surgical technique, assessment of animal well-being, appropriate use of analgesics, and animal physiologic status during all phases of a protocol involving surgery and postoperative care (see Appendix A, Anesthesia, Pain, and Surgery). The individual impact of those factors will vary according to the complexity of procedures involved and the species of animal used. A team approach to a surgical project often increases the likelihood of a successful outcome by providing input from persons with different expert erg(c)-3(o)--3(o)-3(m)-3( )-1w in113 po th

traf c (AORN 2006; Bartley 1993). If it is necessary to use an operating room for other purposes, it is imperative that the room be returned to an appropriate level of hygiene before its use for major survival surgery.

major or minor should be evaluated on a case-by-case basis by the veterinarian and IACUC.

Emergency situations sometimes require immediate surgical attention under less than ideal conditions. For example, if an animal maintained outdoors needs surgical attention, movement to a surgical facility might be impractical or pose an unacceptable risk to the animal. Such situations often require more intensive aftercare and may pose a greater risk of postoperative

Speci c sterilization methods should be selected on the basis of the physical characteristics of the materials to be sterilized (Callahan et al. 1995; Scho eld 1994) and sterilization indicators should be used to validate that materials have been properly sterilized (Berg 1993). Autoclaving and plasma and gas sterilization are effective methods most commonly used to sterilize instruments and materials. Alternative methods, used primarily for rodent surgery, include liquid chemical sterilants and dry heat sterilization. Liquid chemical sterilants should be used with appropriate contact times and instruments should be rinsed with sterile water or saline before use. Bead or dry heat sterilizers are an effective and convenient means of rapidly sterilizing the working surfaces of surgical instruments but care should be taken to ensure that the instrument surfaces have cooled suf ciently before touching animal tissues to minimize the risk of burns. Alcohol is neither a sterilant nor a high-level disinfectant (Rutala 1990) but may be acceptable for some procedures if prolonged contact times are used (Huerkamp 2002).

Intraoperative Monitoring

Careful monitoring and timely attention to problems increase the likelihood of a successful surgical outcome (Kuhlman 2008). Monitoring includes routine evaluation of anesthetic depth and physiologic functions and conditions, such as body temperature, cardiac and respiratory rates and pattern (Flegal et al. 2009), and blood pressure (Kuhlman 2008), and should be appropriately documented. Use of balanced anesthesia, including the addition of an intraoperative analgesic agent, can help minimize physiologic uctuations during surgery. Maintenance of normal body temperature minimizes cardiovascular and respiratory disturbances caused by anesthetic

trained personnel. Particular attention should be given to thermoregulation, cardiovascular and respiratory function, electrolyte and uid balance, and management of postoperative pain or discomfort. Additional care may be warranted, including long-term administration of parenteral uids, analgesics, and other drugs, as well as care of surgical incisions. Appropriate medical records should also be maintained.

After recovery from anesthesia, monitoring is often less intense but should include attention to basic biologic functions of intake and elimination and to behavioral signs of postoperative pain, monitoring top65(t)-3(o)o-1 54 5tc

(all), anorexia (all), rapid or labored respiration (rodents, birds, sh), lack of grooming (mammals and birds), increased aggression (mammals

used in combination with appropriate analgesics and anesthetics to provide balanced anesthesia and to minimize stress associated with perioperative procedures. Neuromuscular blocking agents (e.g., pancuronium) are sometimes used to paralyze skeletal muscles during surgery in which general anesthetics have been administered (Klein 1987); because this paralysis eliminates many signs and reflexes used to assess anesthetic depth, autonomic nervous system changes (e.g., sudden changes in heart rate and blood pressure) can be indicators of pain related to an inadequate depth of anesthesia. It is imperative that any proposed use of neuromuscular blocking drugs be carefully evaluated by the veterinarian and IACUC to ensure the well-being of the animal. Acute stress is believed to be a consequence of paralysis in a conscious state and it is known that humans, if conscious, can experience distress when paralyzed with these drugs (NRC 2008; Van Sluyters and Oberdorfer 1991). If paralyzing agents are to be used, the appropriate amount of anesthetic should first be defined on the basis of results of a similar procedure using the anesthetic without a blocking agent (NRC 2003, 2008, 2009a).

EUTHANASIA

Euthanasia is the act of humanely killing animals by methods that induce rapid unconsciousness and death without pain or distress. Unless a deviation is justified for scientific or medical reasons, methods should be consistent with the AVMA Guidelines on Euthanasia (AVMA 2007 or later editions). In evaluating the appropriateness of methods, some of the criteria that should be considered are ability to induce loss of consciousness and death with no or only momentary pain, distress, or anxiety; reliability; irreversibility; time required to induce unconsciousness; appropriateness for the species and age of the animal; compatibility with research objectives; and the safety of and emotional effect on personnel.

Euthanasia may be planned and necessary at the end of a protocol or as a means to relieve pain or distress that cannot be alleviated by analgesics, sedatives, or other treatments. Criteria for euthanasia include protocol-specific endpoints (such as degree of a physical or behavioral deficit or tumor size) that will enable a prompt decision by the veterinarian and the investigator to ensure that the endpoint is humane and, whenever possible, the scientific objective of the protocol is achieved (see Chapter 2).

Standardized methods of euthanasia that are predictable and controllable should be developed and approved by the AV and IACUC. Euthanasia should be carried out in a manner that avoids animal distress. Automated systems for controlled and staged delivery of inhalants may offer advantages for species killed frequently or in large numbers, such as rodents (McIntyre et al. 2007). Special consideration should be given to euthanasia of fetuses

and larval life forms depending on species and gestational age (Artwohl et al. 2006).

The selection of specific agents and methods for euthanasia will depend on the species involved, the animal's age, and the objectives of the protocol. Generally, chemical agents (e.g., barbiturates, nonexplosive inhalant anesthetics) are preferable to physical methods (e.g., cervical dislocation, decapitation, use of a penetrating captive bolt); however, scientific considerations may preclude the use of chemical agents for some protocols.

Although carbon dioxide (CO_2) is a commonly used method for rodent euthanasia, there is ongoing controversy about its aversive characteristics as an inhalant euthanasia agent. This is an area of active research (Conlee et al. 2005; Danneman et al. 1997; Hackbarth et al. 2000; Kirkden et al. 2008; Leach et al. 2002; Niel et al. 2008) and further study is needed to optimize the methods for CO_2 euthanasia in rodents (Hawkins et al. 2006). The acceptability of CO_2 as a euthanasia agent for small rodents should be evaluated as new data become available. Furthermore, because neonatal rodents are resistant to the hypoxia-inducing effects of CO_2 and require longer exposure times to the agent (Artwohl et al. 2006), alternative methods should be considered (e.g., injection with chemical agents, cervical dislocation, or decapitation; Klaunberg et al. 2004; Pritchett-Corning 2009).

It is essential that euthanasia be performed by personnel skilled in methods for the species in question and in a professional and compassionate manner. Special attention is required to ensure proficiency when a physical method of euthanasia is used. Death must be confirmed by personnel trained to recognize cessation of vital signs in the species being euthanized. A secondary method of euthanasia (e.g., thoracotomy or exsanguination) can be also used to ensure death. All methods of euthanasia should be reviewed and approved by the veterinarian and IACUC.

Euthanizing animals is psychologically difficult for some animal care, veterinary, and research personnel, particularly if they perform euthanasia repetitively or are emotionally attached to the animals being euthanized (Arluke 1990; NRC 2008; Rollin 1986; Wolfe 1985). When delegating euthanasia responsibilities, supervisors should be sensitive to this issue.

REFERENCES

Anderson LC. 2007. Institutional and IACUC responsibilities for animal care and use education and training programs. ILAR J 48:90-95.

ACLAM [American College of Laboratory Animal Medicine]. 2001. Position Statement on Rodent Surgery. Available at www.aclam.org/education/guidelines/position_rodentsurgery.html; accessed January 7, 2010.

AORN [Association of Operating Room Nurses]. 2006. Recommended practices for traffic patterns in the perioperative practice setting. AORN J 83:681-686.

Arluke A. 1990. Uneasiness among laboratory technicians. Lab Anim 19:20-39.

Arndt SS, Lohavech D, van't Klooster J, Ohl F. 2010. Co-species housing in mice and rats: Ef

Carstens E, Moberg GP. 2000. Recognizing pain and distress in laboratory animals. ILAR J 41:62-71.

Chamberlain GV, Houang E. 1984. Trial of the use of masks in gynecological operating theatre. Ann R Coll Surg Engl 66:432-433.

Kagira JM, Ngotho M, Thuita JK, Maina NW, Hau J. 2007. Hematological changes in vervet monkeys (Chlorocebus aethiops) during eight months' adaptation to captivity. Am J Primatol 69:1053-1063.

Karas AZ. 2002. Postoperative analgesia in the laboratory mouse (Mus musculus Lab Anim (NY) 31:49-52.

Karas A, Danneman P, Cadillac J. 2008. Strategies for assessing and minimizing pain. In: Fish R, Brown M, Danneman P, Karas A, eds. Anesthesia and Analgesia in Laboratory Animals. San Diego: Academic Press. p 195-218.

Kirkden RD, Niel L, Stewart SA, Weary DM. 2008. Gas killing of rats: The effect of supplemental oxygen on aversion to carbon dioxide. Anim Welf 17:79-87.

Klaunberg BA, O'Malley J, Clark T, Davis JA. 2004. Euthanasia of mouse fetuses and neonates. Contemp Top Lab Anim Sci 43:29-34.

Klein L. 1987. Neuromuscular blocking agents. In: Short CE, ed. Principles and Practice of Veterinary Anesthesia. Baltimore: Williams and Wilkins. p 134-153.

Klement P, del Nido PJ, Mickleborough L, MacKay C, Klement G, Wilson GJ. 1987. Techniques and postoperative management for successful cardiopulmonary bypass and open-heart surgery in dogs. JAVMA 190:869-874.

Kohn DF, Martin TE, Foley PL, Morris TH, Swindle MM, Vogler GA, Wixson SK. 2007. Public statement: Guidelines for the assessment and management of pain in rodents and rabbits. JAALAS 46:97-108.

Kona-Boun JJ, Silim A, Troncy E. 2005. Immunologic aspects of veterinary anesthesia and analgesia. JAVMA 226:355-363.

Kuhlman SM. 2008. Principles of anesthesia monitoring: Introduction. J Invest Surg 21:161-162.

Landi MS, Kreider JW, Lang CM, Bullock LP. 1982. Effects of shipping on the immune function in mice. Am J Vet Res 43:1654-1657.

Leach MC, Bowell VA, Allan TF, Morton DB. 2002. Aversion to gaseous euthanasia agents in rats and mice. Comp Med 52:249-257.

Lerche NW, Yee JL, Capuano SV, Flynn JL. 2008. New approaches to tuberculosis surveillance in nonhuman primates. ILAR J 49:170-178.

Lovaglio J, Lawson PT. 1995. A draping method for improving aseptic technique in rodent surgery. Lab Anim (NY) 24:28-31.

Maggio-Price L, Shows D, Waggie K, Burich A, Zeng W, Escobar S, Morrissey P, Viney JL. 2002. Helicobacter bilis

Morton CM, Reid J, Scott EM, Holton LL, Nolan AM. 2005. Application of a scaling model to establish and validate an interval level pain scale for assessment of acute pain in dogs. Am J Vet Res 66:2154-2166.

Murphey-Corb M, Martin LN, Rangan SRS, Baskin GB, Gormus BJ, Wolf RH, Andes WA, West M, Montelaro RC. 1986. Isolation of an HTLV-III-related retrovirus from macaques with simian AIDS and its possible origin in asymptomatic mangabeys. Nature 321:435-437.

Murphy BL, Maynard JE, Krushak DH, Fields RM. 1971. Occurrence of a carrier state herpesvirus tamarinus in marmosets. Appl Microbiol 21:50-52.

Murrell JC, Johnson CB. 2006. Neurophysiological techniques to assess pain in animals. J Vet Pharmacol Ther 29:325-335.

Nicklas W, Kraft V, Meyer B. 1993. Contamination of transplantable tumors, cell lines, W (e)-3(.1233(e)-

Pritchett-Corning KR. 2009. Euthanasia of neonatal rats with carbon dioxide. JAALAS 48:23-27.

Pritchett-Corning KR, Chang FT, Festing MF. 2009. Breeding and housing laboratory rats and mice in the same room does not affect the growth or reproduction of either species. JAALAS 48:492-498.

Renquist D. 1990. Outbreak of simian hemorrhagic fever. J Med Primatol 19:77-79.

Ritter MA, Marmion P. 1987. The exogenous sources and controls of microorganisms in the operating room. Orthop Nurs 7:23-28.

Roberts RA, Andrews K. 2008. Nonhuman primate quarantine: Its evolution and practice. ILAR J 49:145-156.

Robertshaw D. 2004. Temperature regulation and the thermal environment. In: Duke's Physiol ogy of Domestic Animals, 12th ed. WO Reese, ed. Ithaca, NY: Cornell University Press.

Robinson V, Anderson S, Carver JFA, Francis RJ, Hubrecht R, Jenkins E, Mathers KE, Raymond R, Rosewell I, Wallace J, Wells DJ. 2003. Re nement and reduction in production of genetically modi ed mice. Sixth report of the BVAAWF/FRAME/RSPCA/UFAW Joint Working Group on Re nement. Lab Anim 37(Suppl 1):S1-S51. Available at www.lal.org.uk/pdf les/Transgenic.pdf; accessed January 7, 2010.

Rollin B. 1986. Euthanasia and moral stress. In: DeBellis R, ed. Loss, Grief and Care. Bing hamton NY: Haworth Press.

Rose JD. 2002. The neurobehavioral nature of shes and the question of awareness and pain. Rev Fish Sci 10:1-38.

Roughan JV, Flecknell PA. 2000. Behavioural effects of laparotomy and analgesic effects of ketoprofen and carprofen in rats. Pain 90:65-74.

Roughan JV, Flecknell PA. 2003. Evaluation of a short duration behaviour-based post-operative pain scoring system in rats. Eur J Pain 7:397-406.

Roughan JV, Flecknell PA. 2004. Behaviour-based assessment of the duration of laparotomy-induced abdominal pain and the analgesic effects of carprofen and buprenorphine in rats. Behav Pharmacol 15:461-472.

Rutala WA. 1990. APIC guideline for selection and use of disinfectants. Am J Infect Contr 18:99-117.

Sanhouri AA, Jones RS, Dobson H. 1989. The effects of different types of transportation on plasma cortisol and testosterone concentrations in male goats. Br Vet J 145:446-450.

Scho eld JC. 1994. Principles of aseptic technique. In: Bennett BT, Brown MJ, Scho eld JC, eds. Essentials for Animal Research: A Primer for Research Personnel. Washington: Na tional Agricultural Library. p 59-77.

Schonholtz GJ. 1976. Maintenance of aseptic barriers in the conventional operating room. J Bone Joint Surg 58:439-445.

Schrama JW, van der Hel W, Gorssen J, Henken AM, Verstegen MW, Noordhuizen JP. 1996. Required thermal thresholds during transport of animals. Vet Q 18(3):90-95.

Sherwin CM. 2001. Can invertebrates suffer? Or, how robust is argument by analogy? Anim Welf 10:103-118.

Slattum MM, Maggio-Price L, DiGiacomo RF, Russell RG. 1991. Infusion-related sepsis in dogs undergoing acute cardiopulmonary surgery. Lab Anim Sci 41:146-150.

Sneddon LU. 2006. Ethics and welfare: Pain perception in sh. Bull Eur Assoc Fish Pathol 26:6-10.

Spinelli J. 1990. Preventing suffering in laboratory animals. In: Rollin B, Kesel M, eds. The Experimental Animal in Biomedical Research, vol I: A Survey of Scienti c and Ethical Issues for Investigators. Boca Raton, FL: CRC Press. p 231-242.

Stevens CA, Dey ND. 2007. A program for simulated rodent surgical training. Lab Anim (NY) 36:25-31.

Stoskopf MK. 1994. Pain and analgesia in birds, reptiles, amphibians, and sh. Invest Oph thalmol Vis Sci 35:775-780.
Suckow MA, Doerning BJ. 2007. Assessment of Veterinary Care. In: Silverman J, Suckow MA, Murthy S, eds. The IACUC Handbook, 2nd ed. Boca Raton, FL: CRC Press.
Tuli JS, Smith JA, Morton DB. 1995. Stress measurements in mice after transportation. Lab Anim 29:132-138.
UFAW [Universities Federation for Animal Welfare]. 1989. Surgical procedures. In: Guidelines on the Care of Laboratory Animals and Their Use for Scienti c Purposes III. London. p 3-15.
USDA [US Department of Agriculture]. 1985. 9 CFR 1A. (Title 9, Chapter 1, Subchapter A): Animal Welfare. Available at http://ecfr.gpoaccess.gov/cgi/t/text/text-idx?sid=8314313bd 7adf2c9f1964e2d82a88d92andc=ecfrandtpl=/ecfrbrowse/Title09/9cfrv1_02.tpl; accessed January 14, 2010.
USDA. 1997. APHIS Policy #3, “Veterinary Care” (July 17, 1997). Available at www.aphis. usda.gov/animal_welfare/downloads/policy/policy3.pdf; accessed January 9, 2010.
Valverde A, Gunkel CI. 2005. Pain management in horses and farm animals. J Vet Emerg Crit Care 15:295-307.
Van Sluyters RC, Oberdorfer MD, eds. 1991. Preparation and Maintenance of Higher Mammals During Neuroscience Experiments. Report of a National Institutes of Health Workshop. NIH No. 91-3207. Bethesda, MD.
Viñuela-Fernández I, Jones E, Welsh EM, Fleetwood-Walker SM. 2007. Pain mechanisms and their implication for the management of pain in farm and companion animals. Vet J 174:227-239.
Waynforth HB. 1980. Experimental and Surgical Technique in the Rat. London: Academic Press.
Waynforth HB. 1987. Standards of surgery for experimental animals. In: Tuffery AA, ed. Labo ratory Animals: An Introduction for New Experimenters. Chichester: Wiley-Interscience. p 311-312.
Whyte W. 1988. The role of clothing and drapes in the operating room. J Hosp Infect 11(Suppl C):2-17.
Wol e TL. 1985. Laboratory animal technicians: Their role in stress reduction and human-companion animal bonding. Vet Clin N Am Small Anim Pract 15:449-454.

Building materials for animal facilities should be selected to facilitate ef cient and hygienic operation. Durable, moisture- and vermin-proof, re-resistant, seamless materials are most desirable for interior surfaces, which should be highly resistant to the effects of cleaning agents, scrubbing, high-pressure sprays, and impact. Paints and glazes should be nontoxic if used on surfaces with which animals will have direct contact. In the construction of outdoor facilities, consideration should be given to surfaces that withstand the elements and can be easily maintained.

Location

Quality animal management and human comfort and health protection require separation of animal facilities from personnel areas, such as of ces and conference rooms. Separation can be accomplished by having the animal quarters in a separate building, wing, oor, or room. Careful planning should make it possible to place animal housing areas next to or near research laboratories but separated from them by barriers, such as entry locks, corridors, or oors. Additional considerations include the impact of noise and vibration generated from within the facility and from surrounding areas of the building, as well as security of the facility.

Animals should be housed in facilities dedicated to or assigned for that purpose, not in laboratories merely for convenience. If animals must be maintained in a laboratory to satisfy the scienti c aims of a protocol, that space should be appropriate to house and care for the animals and its use limited to the period during which it is required. If needed, measures should be taken to minimize occupational hazards related to exposure to animals both in the research area and during transport to and from the area.

Centralization Versus Decentralization

In a physically centralized animal facility, support, care, and use areas are adjacent to the animal housing space. Decentralized animal housing and use occur in space that is not solely dedicated to animal care or support or is physically separated from the support areas and animal care personnel. Centralization often reduces operating costs, providing a more ef cient ow of animal care supplies, equipment, and personnel; more ef cient use of environmental controls; and less duplication of support services. Centralization reduces the needs for transporting animals between housing and study sites, thereby minimizing the risks of transport stress and exposure to disease agents; affords greater security by providing the opportunity to control facility access; and increases the ease of monitoring staff and animals.

Decentralized animal facilities generally cost more to construct because of the requirement for specialized environmental systems and

controls in multiple sites. Duplicate equipment (e.g., cage washers) may be needed, or soiled materials may need to be moved distances for processing. But decentralization may be preferred for certain specialized research services such as imaging, quarantine, and proximity to research facilities, or for biosecurity reasons. Decentralization may be necessary to accommodate large or complex equipment, such as magnetic resonance imaging, or to permit space sharing by users from multiple facilities or institutions. The opportunity for exposure to disease agents is much greater in these situations and special consideration should be given to biosecurity, including transportation to and from the site, quarantine before or after use of the specialized research area, and environmental and equipment decontamination.

The decisions leading to selection of physically centralized versus decentralized animal facilities should be made early and carefully and should involve all stakeholders (NRC 1996; Ruys 1991).

FUNCTIONAL AREAS

Professional judgment should be exercised in the development of a practical, functional, and ef cient physical plant for animal care and use. The size, nature, and intensity of an institutional Program (see Chapter 2) will determine the speci c facility and support functions needed. In facilities that are small, maintain few animals, or maintain animals under special conditions—such as facilities used exclusively for housing gnotobiotic or speci c pathogen-free (SPF) colonies or animals in runs, pens, or outdoor housing—some functional areas listed below may be unnecessary or may be included in a multipurpose area.

Space is required for the following:

- animal housing, care, and sanitation
- receipt, quarantine, separation, and/or rederivation of animals
- separation of species or isolation of individual projects when necessary
- storage.

Most multipurpose animal facilities may also include the following:

- specialized laboratories or space contiguous with or near animal housing areas for such activities as surgery, intensive care, necropsy, irradiation, preparation of special diets, experimental procedures, behavioral testing, imaging, clinical treatment, and diagnostic laboratory procedures

- containment facilities or equipment, if hazardous biologic, physical, or chemical agents are to be used
- barrier facilities for housing of SPF rodents, especially valuable genetically modi ed animals, or irreplaceable animal models

becoming gouged, cracked, or pitted. Depending on their use, oors should be monolithic or have a minimal number of joints. Some materials that have

Power and Lighting

The electrical system should be safe and provide appropriate lighting, a suf cient number of power outlets, and suitable amperage for specialized equipment. In the event of power failure, an alternative or emergency power supply should be available to maintain critical services (e.g., the HVAC system, ventilated caging systems [Huerkamp et al. 2003], or life support systems for aquatic species) or support functions (e.g., freezers and isolators) in animal rooms, operating suites, and other essential areas. Consideration should be given to out tting movable equipment for which uninterrupted power is essential (e.g., ventilated racks), with twist-lock plugs to prevent accidental removal from the power supply.

Light xtures, timers, switches, and outlets should be properly sealed to prevent vermin access. Recessed energy-ef cient uorescent lights are commonly used in animal facilities. Spectral quality of lights may be important for some species when maintained in the laboratory; in these cases full spectrum lamps may be appropriate. A time-controlled lighting system should be used to ensure a uniform diurnal lighting cycle. Override systems should be equipped with an automatic timeout or a warning light

storage areas. Refrigerated storage, separated from other cold storage, is

SPECIAL FACILITIES

Surgery

The design of a surgical facility should accommodate the species to be operated on and the complexity of the procedures to be performed (Hessler 1991; see also Appendix A, Design and Construction of Animal Facilities). The facility, including that used for rodents, by necessity becomes larger and more complex as the number and size of animals or the complexity of procedures increase. For instance, a larger facility may be required to accommodate procedures on agricultural species, large surgical teams, imaging devices, robotic surgical systems, and/or laparoscopic equipment towers. Surgical facilities for agricultural species may additionally require oor drains, special restraint devices, and hydraulic operating tables.

For most survival surgery performed on rodents and other small species such as aquatics and birds, an animal procedure laboratory is recommended; the space should be dedicated to surgery and related activities when used for this purpose, and managed to minimize contamination from other activities conducted in the room at other times. The association of surgical facilities with diagnostic laboratories, imaging facilities, animal housing, staff of ces, and so on should be considered in the overall context of the complexity of the surgical program. Centralized surgical facilities are cost-effective in equipment, conservation of space and personnel resources, and reduced transit of animals. They also enable enhanced personnel safety and professional oversight of both facilities and procedures.

For most surgical programs, functional components of aseptic surgery include surgical support, animal preparation, surgeon's scrub, operating room, and postoperative recovery. The areas that support those functions should be designed to minimize traf c ow and separate the related nonsurgical activities from the surgical procedure in the operating room. The separation is best achieved by physical barriers (AORN 1993) but may also be achieved by distance between areas or by the timing of appropriate cleaning and disinfection between activities.

Surgical facilities should be suf ciently separate from other areas to minimize unnecessary traf c and decrease the potential for contamination (Humphreys 1993). The number of personnel and their level of activity have been shown to be directly related to the level of bacterial contamination and the incidence of postoperative wound infection (Fitzgerald 1979). Traf c in the operating room can be reduced by the installation of an observation window, a communication system (such as an intercom system), and judicious location of doors.

Control of contamination and ease of cleaning should be key considerations in the design of a surgical facility. The interior surfaces should be

and rats, especially valuable genetically engineered animals, and SPF animals of other species.

Barrier facilities typically incorporate airlock or special entries (e.g., air or wet showers) for staff and supplies. Staff generally wear dedicated clothing and footwear, or freshly laundered, sterile, or disposable outer garments such as gowns, head and shoe covers, gloves, and sometimes face masks prior to entry. Consumables, such as feed or bedding, that may harbor infectious agents are autoclaved or are gamma-irradiated by the supplier and surface decontaminated on entry. Drinking water may be autoclaved or subject to specialized treatment (e.g., reverse osmosis ltration) to remove infectious agents. Caging and other materials with which the animals have direct contact may be sterilized after washing before reuse. Strict operational procedures are frequently established to preclude intermingling of clean and soiled supplies and personnel groups, depending on work function. Only animals of de ned health status are received into the barrier, and once they leave they are prohibited from reentering without retesting. Personnel entry is restricted and those with access are appropriately trained in procedures that minimize the introduction of contaminants.

Engineering features may include high-level ltration of supply air (e.g., HEPA or 95% ef cient lters), pressurization of the barrier with respect to surrounding areas, and directional air ow from clean to potentially contaminated areas. Specialized equipment augmenting the barrier may include isolator cages, individually ventilated cages, and animal changing stations.

Detailed information on barrier design, construction, and operations has been recently published (Hessler 2008; Lipman 2006, 2008).

Imaging

In vivo imaging offers noninvasive methods for evaluating structure

contamination between groups of animals, different animal species, or between animals and humans (if the device is used for both animal and human subjects) is possible because these devices may be difficult to sanitize (Klaunberg and Davis 2008; Lipman 2006). If the imaging resource is located outside the animal facility, appropriate transportation methods and routes should be developed to avoid inappropriate exposure of humans to animals in transit. If possible, animals should not be moved past offices, lunch rooms, or public areas where people are likely to be present.

As imaging may require the subject to be immobile, often for extended time periods during image acquisition, provisions should be made for delivery of anesthetics and carrier gas, the scavenging of waste anesthetic gas, and adequate animal monitoring (Balaban and Hampshire 2001). Remote storage of gas tanks is generally required in facilities where magnetic resonance (MR) scanners are used as the magnetic field requires ferrous materials to be kept a safe distance away from the magnet. Site selection of MR scanners requires special attention because of their weight, the fringe field generated (especially from unshielded magnets), and the impact of ferrous elements of the building structure or its components, especially those that are not static (e.g., elevators), as they may affect field homogeneity. Most MR scanners are superconducting and require the use of cryogens. Because cryogen boil-off can lead to asphyxiation of both personnel and animals, rooms with MR scanners or in which cryogen gases are stored must be equipped with oxygen sensors and a method for increasing room ventilation to exhaust inert gases during cryogen filling (Klaunberg and Davis 2008).

Many imaging devices, especially those designed for small animals, are self-contained and require no special physical plant considerations. Provisions should be made to locate the operating console away from imaging devices that emit ionizing or magnetic radiation. Imaging devices with components that are difficult to sanitize should be covered with a disposable or sanitizable material when not in use.

Whole Body Irradiation

Total body irradiation of small laboratory animals may be accomplished using devices that emit either gamma- or X-rays. Devices are usually self-shielded and, because of the weight of the shielding material, may require special site considerations. Devices with gamma-emitting sources are subject to regulations that require adherence to specific security, monitoring, and personnel clearance requirements (Nuclear Regulatory Commission 2008). The site selected for irradiators should also take into account whether they are to be used for animals and biologics, as well as the source and microbial status of the animals to be irradiated. Locating them in the animal facility may require access for personnel who would normally not

require it or may necessitate bringing animals into a facility where they are not normally housed.

Hazardous Agent Containment

The goal of containment is to "reduce or eliminate exposure of laboratory workers, other persons, and the outside environment to potential hazardous

wo sintnas

ASHRAE [American Society of Heating, Refrigeration, and Air Conditioning Engineers]. 2007a. Chapter 22: Environmental control for animals and plants. In 2007 ASHRAE Handbook: Fundamentals, I-P ed. Atlanta.

ASHRAE. 2007b. Chapter 47: Sound and vibration control. In 2007 ASHRAE Handbook: HVAC Applications. Atlanta.

ASHRAE. 2007c. Chapter 45: Control of gaseous indoor air contaminants. In 2007 ASHRAE Handbook: HVAC Applications. Atlanta.

Heffner HE, Heffner RS. 2007. Hearing ranges of laboratory animals. JAALAS 46:20-22.
Hessler JR. 1991. Facilities to support research. In: Ruys T, ed. Handbook of Facility Planning, vol 2: Laboratory Animal Facilities. New York: Van Nostrand. p 34-55.
Hessler JR. 2008. Barrier housing for rodents. In: Hessler J, Lehner N, eds. Planning and Designing Animal Research Facilities. Amsterdam: Academic Press. p 335-345.
Holton J, Ridgway GL. 1993. Commissioning operating theatres. J Hosp Infect 23:153-160.
Huerkamp MJ, Thompson WD, Lehner NDM. 2003. Failed air supply to individually ventilated caging system causes acute hypoxia and mortality of rats. Contemp Top Lab Anim Sci 42:44-45.
Humphreys H. 1993. Infection control and the design of a new operating theatre suite. J Hosp Infect 23:61-70.
Jacobs GH, Fenwick JA, Williams GA. 2001. Cone-based vision of rats for ultraviolet and visible lights. J Exp Biol 204:2439-2446.
Klaunberg BA, Davis JA. 2008. Considerations for laboratory animal imaging center design and setup. ILAR J 49:4-16.
Knecht CD, Allen AR, Williams DJ, Johnson JH. 1981. Fundamental Techniques in Veterinary Surgery, 2nd ed. Philadelphia: WB Saunders.
Kowalski WJ, Bahn eth WP, Carey DD. 2002. Engineering control of airborne disease transmission in animal laboratories. Contemp Top Lab Anim Sci 41:9-(1)-3(a)-(1)-3(:mh-3(i)-3( )20 Td (-(.)-3( )

TmIdaadgumRtoad2.25 -1.25tnary

Ross S, Schapiro S, Hau J, Lukas K. 2009. Space use as an indicator of enclosure appropriate ness: A novel measure of captive animal welfare. Appl Anim Behav Sci 121:42-50.

Addendum

Guide for the Care and Use of Laboratory Animals Eighth Edition

ADDENDUM: LIST OF EDITORIAL CHANGES FROM THE PREPUBLICATION VERSION[1]

1. Page 22. Original sentence: “Personnel training should include information on laboratory animal allergies, preventive control mea sures and proper techniques for working with animals (Gordon et at. 1997; Schweitzer et al. 2003; Thulin et al. 2002).” “…early recognition and reporting of allergy symptoms” was added to clarify the guidance and re ect the cited references.

2. Page 31. Original sentence: “They should therefore be used, when available, for all animal-related procedures (NIH 2008; USDA 1997b).” The NIH reference was removed because it applies only to the NIH intramural animal research program.

3. Page 32. Original sentence: “Principal investigators conducting eld research should be knowledgeable of relevant zoonotic diseases, associated safety issues, and, when working in an international environment, any local laws or regulations that apply.” Compliance with laws and regulations applies to eld investigations irrespective of location, so “when working in an international environment” and “local” were deleted.

[1]Page numbers re ect placement of revisions in this report.

10. Page 107. Original sentence: “The Centers for Disease Control and Prevention enforces regulations to prevent the introduction, transmission, or spread of communicable diseases and regulate the importation of any animal or animal product capable of carrying a zoonotic disease.” Because USDA also has jurisdiction over imports, it was added to the sentence.

11. Page 107. Original sentence: “The US Fish3(e)-3(g)-3(e8-)-150(-3(e)-3( )-2

Appendices

APPENDIX
A

Additional Selected References

SUBJECT MATTER

USE OF LABORATORY ANIMALS
Alternatives
Ethics and Welfare
Experimental Design and Statistics
Research and Testing Methodology

PROGRAM MANAGEMENT
General References
Laws, Regulations, and Policies
Education
Monitoring the Care and Use of Animals
Occupational Health and Safety

ENVIRONMENT, HOUSING, AND MANAGEMENT
General References
Environmental Enrichment
Genetics and Genetically Modi ed Animals
Species-Speci c References—Environment, Housing, and Management
  Agricultural Animals
  Amphibians, Reptiles, and Fish
  Birds
  Cats and Dogs
  Exotic, Wild, and Zoo Animals

Nonhuman Primates
Rodents and Rabbits
Other Animals

VETERINARY CARE
Transportation
Anesthesia, Pain, and Surgery
Disease Surveillance, Diagnosis, and Treatment
Pathology, Clinical Pathology, and Parasitology
Species-Speci c References—Veterinary Care
Agricultural Animals
Amphibians, Reptiles, and Fish
Birds
Cats and Dogs
Exotic, Wild, and Zoo Animals
Nonhuman Primates
Rodents and Rabbits

Re nement, reduction, and replacement of animal use for regulatory testing: Future im provements and implementation within the regulatory framework. 2002. Richmond J. ILAR J 43:S63-S68.

The Role of the Interagency Coordinating Committee on the Validation of Alternative Methods (ICCVAM) in the Evaluation of New Toxicological Testing Methods. 2000. Stokes WS, Hill RN. In: Proceedings of the 3rd World Congress on Alternatives and Animal Use in the Life Sciences, Bologna, Italy, 1999. New York: Elsevier.

Ethics and Welfare

An additional “R”: Remembering the animals. 2002. Iliff SA. ILAR J 43:38-47.

Animal Liberation, 2nd ed. 1990. **Singer P. New York: New York Review Book (distributed by** Random House).

Humane endpoints for laboratory animals used in toxicity testing. 2000. Stokes WS. In: Proceedings of the 3rd World Congress on Alternatives and Animal Use in the Life Sciences, Bologna, Italy, 1999. New York: Elsevier.

Humane endpoints in animal experiments for biomedical research: Proceedings of the International Conference, 22-25 November 1998, Zeist, the Netherlands.

Immunization Procedures and Adjuvant Products. 2005. ILAR J 46(3).

Adjuvants and antibody production: Dispelling the myths associated with Freund's complete and other adjuvants. 2005. Stills JF Jr. ILAR J 46:280-293.

Advances in monoclonal antibody technology: Genetic engineering of mice, cells, and immunoglobulins. 2005. Peterson NC. ILAR J 46:314-319.

Applications and optimization of immunization procedures. 2005. Schunk MK, Macallum GE. ILAR J 46:241-257.

Monoclonal versus polyclonal antibodies: Distinguishing characteristics, applications, and information. 2005. Lip 5.i Li.d L.iR3(i)-34-3(.)c-8-Trudel-3(i)-1LJ3(i)-34-3(.)c-8-Weis-Garcia(.) 46:258-268.

Using polyclonal and monoclonal antibodies in regulatory testing of biological products.

Future improvements and implementation of animal care practices within the animal test ing regulatory environment. 2002. Guittin P, Decelle T. ILAR J 43:S80-S84.
Incorporating the 3Rs into regulatory scienti c practices. 2002. Sterling S, Rispin A. ILAR J 43:S18-S20.
Possibilities for re nement and reduction: Future improvements within regulatory testing. 2002. Stephens ML, Conlee K, Alvino G, Rowan A. ILAR J 43:S74-S79.

Introduction: Disaster planning and management: A practicum. 2010. Bayne KA. ILAR J 51:101-103.
IACUC considerations: You have a disaster plan but are you really prepared? 2010. Wing eld WE, Rollin BE, Bowen RA. ILAR J 51:164-170.
Management of rodent viral disease outbreaks: One institution's (r)evolution. 2010. Smith AL. ILAR J 51:127-137.
Tropical storm and hurricane recovery and preparedness strategies. 2010. Goodwin BS Jr, Donaho JC. ILAR J 51:104-119.
Veri cation of poultry carcass composting research through application during actual avian in uenza outbreaks. 2010. Flory GA, Peer RW. ILAR J 51:149-157.
Wild re evacuation: Outrunning the witch's curse—One animal center's experience. 2010. Arms MM, Van Zante JD. ILAR J 51:158-163.
Essentials for Animal Research: A Primer for Research Personnel. 1994. Bennett BT, Brown MJ, Scho eld JC. Beltsville MD: National Agricultural Library.
Infectious Disease Research in the Age of Biodefense. ILAR J 46(1).
Administrative issues related to infectious disease research in the age of bioterrorism. 2005. Jaax J. ILAR J 46:8-14.
Public response to infectious disease research: The UC Davis experience. 2005. Fell AH, Bailey PJ. ILAR J 46:66-72.
Laboratory security and emergency response guidance for laboratories working with select agents. Richmond JY, Nesby-O'Dell SL. 2002. MMWR Recomm Rep 51:1-8.
Management of Laboratory Animal Care and Use Programs. 2002. Suckow MA, Douglas FA, Weichbrod RH, eds. Boca Raton FL: CRC Press.
Use of Laboratory Animals in Biomedical and Behavioral Research. 1988. National Research Council and Institute of Medicine. Washington: National Academy Press.
Using site assessment and risk analysis to plan and build disaster-resistant programs and facili ties. 2003. Vogelweid CM, Hill JB, Shea RA, Truby SJ, Schantz LD. Lab Anim 32:40-44.

Laws, Regulations, and Policies

Animal Care and Use: Policy Issues in the 1990s. National Institutes of Health/Of ce for Pro tection from Research Risks (NIH/OPRR). 1989. Proceedings of NIH/OPRR Conference, Bethesda MD.
Animal Care Policy Manual. APHIS [Animal and Plant Health Inspection Service], USDA. Available at www.aphis.usda.gov/animal_welfare/policy.shtml; accessed May 15, 2010.
Animal Care Resource Guide: Research Facility Inspection Guide. APHIS, USDA. Available at www.aphis.usda.gov/animal_welfare/rig.shtml; accessed May 15, 2010.
Animal Law Section. National Association for Biomedical Research. Available at http://www.nabranimallaw.org/; accessed October 2, 2010.
Animals and Their Legal Rights. 1985. Washington: Animal Welfare Institute.
APS [American Physiological Society] Guiding Principles for Research Involving Animals and Human Beings. Available at www.the-aps.org/publications/journals/guide.htm; accessed January 24, 2010.

US laws and norms related to laboratory animal research. 1999. VandeBerg JL, Williams-Blangero S, Wol e TL. ILAR J 40:34-37.

Education

AALAS [American Association for Laboratory Animal Science] Technician Training and Certi cation. Available at www.aalas.org/certi cation/tech_cert.aspx; accessed May 21, 2010.

CALAS [Canadian Association of Laboratory Animal Science] Registry. Available at www.calas-acsal.org/index.php?option=com_content&task=view&id=24&Itemid=106; accessed August 10, 2010.

Clinical Textbook for Veterinary Technicians, 5th ed. 2002. McCurnin DM, Bassert JM, eds. Philadelphia: WB Saunders.

Education and Training in the Care and Use of Laboratory Animals: A Guide for Developing Institutional Programs. 1991. National Research Council. Washington: National Academy Press.

FELASA recommendations for the accreditation of laboratory animal science education and training. 2002. Nevalainen T, Blom HJ, Guaitani A, Hardy P, Howard BR, Vergara P. Lab Anim 36:373-377.

Guidelines on Institutional Animal User Training. Ottawa. National Institutional Animal User Training Program. 1999. Canadian Council on Animal Care. Available at http://ccac.ca/en/CCAC_Programs/ETCC/Intro-coretopics-Web11.htm; accessed May 15, 2010.

National n1.25 Td [(G)-3(u)-3(i)-3(d)-3(e)-Anima1(t)-3(t)-3(a)-3713(n1)-3(i)-3(m)-3(a)-J-WyWeb-J3(e)-3(d)-3

Animal Care and Use Committees Bibliography. 1992. Allen T, Clingerman K. Beltsville MD: US Department of Agriculture, National Agricultural Library (Publication #SRB92-16).

Best practices for animal care committees and animal use oversight. 2002. De Haven R. ILAR J 43(Suppl):S59-S62.

Community representatives and nonscientists on the IACUC: What difference should it make? 1999. Dresser R. ILAR J 40:29-33.

Effective animal care and use committees. 1987. In: Orlans FB, Simmonds RC, Dodds WJ, eds. Laboratory Animal Science, Special Issue, January. Published in collaboration with the Scientists Center for Animal Welfare.

Field studies and the IACUC: Protocol review, oversight, and occupational health and safety considerations. 2007. Laber K, Kennedy BW, Young L. Lab Anim 36:27-33.

Guidelines for the veterinary care of laboratory animals. Report of the FELASA/ECLAM/ESLAV Joint Working Group on Veterinary Care. 2008. Lab Anim 42:1-11. Available at http://la.rsmjournals.com/cgi/reprint/42/1/1; accessed May 21, 2010.

Information Resources for Institutional Animal Care and Use Committees 1985-1999. 1999, rev 2000. AWIC Resources Series No. 7. US Department of Agriculture, National Agricultural Library. Available at www.nal.usda.gov/awic/pubs/IACUC/; accessed January 24, 2010.

Institutional Animal Care and Use Committee Guidebook, 2nd ed. 2002. Applied Research Ethics National Association (ARENA), Of ce of Laboratory Animal Welfare (OLAW), National Institutes of Health. Available at http://grants1.nih.gov/grants/olaw/GuideBook.pdf; accessed January 21, 2010.

Principles and practice in ethical review of animal experiments across Europe: Summary of the report of the FELASA Working Group on Ethical Evaluation of Animal Experiments. 2007. Smith JA, van den Broek FAR, Canto Martorell J, Hackbarth H, Ruksenas O, Zeller W. Lab Anim 41:143-160.

Reference Materials for Members of Animal Care and Use Committees. 1991. Berry DJ. Beltsville MD: US Department of Agriculture, National Agricultural Library (AWIC series #10).

Should IACUCs review scienti c merit of animal research projects? 2004. Mann MD, Prentice ED. Lab Anim (NY) 33:26-31.

Supplementary Resources for Lay Members of Local Ethical Review Processes: Projects Involving Genetically Modi ed Animals. 2004. Lane N, Jennings M. West Sussex UK: RSPCA. Available at www.rspca.org.uk/servlet/BlobServer?blobtable=RSPCABlob&blobcol=urlblob&blobkey=id&blobwhere=1105990223463&blobheader=application/pdf; accessed August 10, 2010.

The IACUC Handbook, 2nd ed. 2007. Silverman J, Suckow MA, Murthy S, eds. Boca Raton FL: CRC Press.

The International Symposium on Regulatory Testing and Animal Welfare: Recommendations on best scienti c practices for animal care committees and animal use oversight. 2002. Richmond J, Fletch A, Van Tongerloo R. ILAR J 43(Suppl):S129-S132.

Occupational Health and Safety

Air quality in an animal facility: Particulates, ammonia, and volatile organic compounds. 1996. Kacergis JB, Jones RB, Reeb CK, Turner WA, Ohman JL, Ardman MR, Paigen B. Am Ind Hyg Assoc J 57:634-640.

Allergy to laboratory mice and rats: A review of its prevention, management, and treatment. 1993. Hunskaar S, Fosse RT. Lab Anim 27:206-221.

An overview of the roles and structure of international high-security veterinary laboratories for infectious animal diseases. 1998. Murray PK. Rev Sci Tech Off Int Epiz 17:426-443.

Animal-associated human infections. 1991. Weinberg AN, Weber DJ. Infect Dis Clin North America 5:1-181.

Animal experimentation in level 4 facilities. 2002. Abraham G, Muschilli J, Middleton D. In: Richmond JY, ed. Anthology of Biosafety: BSL-4 Laboratories. Mundelein IL: American Biological Safety Association. p 343-359.

Animal necropsy in maximum containment. 2002. Wilhelmson CL, Jaax NK, Davis K. In: Richmond JY, ed. Anthology of Biosafety: BSL-4 Laboratories. Mundelein IL: American Biological Safety Association. p 361-402.

Association for Assessment and Accreditation of Laboratory Animal Care (AAALAC) International. Position Statement onCercopithecine herpesvirus1, CHV-1 (Herpesvirus-B). Available at www.aaalac.org/accreditation/positionstatements.cfm; accessed January 24, 2010.

Billions for biodefense: Federal agency biodefense funding, FY2001-FY2005. 2004. Schuler A. Biosecur Bioterror 2:86-96.

Biohazards and Zoonotic Problems of Primate Procurement, Quarantine and Research. 1975. Simmons ML, ed. Cancer Research Safety Monograph Series, vol 2. DHEW Pub. No. (NIH) 76-890. Washington: Department of Health, Education, and Welfare.

Biological Safety Principles and Practices. 2000. Fleming DO, Hunt DL, eds. Washington: ASM Press.

Biosafety in Microbiological and Biomedical Laboratories, 5th ed. 2009. Chosewood CL, Wilson DE, eds. DHHS [Department of Health and Human Services]. Washington: Government Printing Office. Available at http://www.cdc.gov/biosafety/publications/bmbl5/index.htm; accessed July 30, 2010.

Biosafety in the Laboratory: Prudent Practices for Handling and Disposal of Infectious Materials. 1989. National Research Council. Washington: National Academy Press.

Biotechnology Research in an Age of Terrorism. 2004. NRC. Washington: National Academies Press.

Code of Federal Regulations. 1984. Title 40; Part 260, Hazardous Waste Management System: General; Part 261, Identification and Listing of Hazardous Waste; Part 262, Standards Applicable to Generators of Hazardous Waste; Part 263, Standards Applicable to Transporters of Hazardous Waste; Part 264, Standards for Owners and Operators of Hazardous Waste Treatment, Storage, and Disposal Facilities; Part 265, Interim Status Standards for Owners and Operators of Hazardous Waste Treatment, Storage, and Disposal Facilities; and Part 270, EPA-Administered Permit Programs: The Hazardous Waste Permit Program. Washington: Office of the Federal Register. (Part 260 updated April 1994; 261 and 270 updated August 1994; 264 and 265 updated June 1994; 262 and 263 updated 1993)

Evaluation of individually ventilated cage systems for laboratory rodents: Occupational health aspects. 2001. Renstrom A, Bjoring G, Hoglund AU. Lab Anim 35:42-50.

Industrial Biocides. 1988. Payne KR, ed. New York: Wiley.

Infectious Disease Research in the Age of Biodefense. ILAR J 46(1).

Issues related to the use of animals in biocontainment research facilities. Copps J. 2005. ILAR J 46:34-43.

Sel-30(t)-3(h)-3(e-1.25 Td [(I)-3(s)-3(s)-3(u)-3(e)-3(s)-3( )-29(r)-3(e)-3(C)-3()-3(s3((h)-3( )--3(9)-3(4)-3 rd -4 PalLara03(s)-3(t)-3(r)-3(i)-3(a)-3seaa03(s)-3(t)-3(r)-3(i)-3 RE(.)-3( )-30(Pe)-3A, Bjorin3(e)-3(s)w

Laboratory Animals: An Introduction for New Experimenters. 1987. Tuffery AA, ed. Chichester: Wiley Interscience.
Managing the Laboratory Animal Facility, 2nd ed. 2009. Silverman J, ed. Boca Raton FL: CRC Press.
Microbiological Aspects of Bio lms and Drinking Water. 2000. Percival SL, Walker JT, Hunter PR. Boca Raton FL: CRC Press.
Pheromones and Reproduction in Mammals. 1983. Vandenbergh JG, ed. New York: Academic Press.
Practical Animal Handling. 1991. Anderson RS, Edney ATB, eds. Elmsford NY: Pergamon.
Recent advances in sterilization. 1998. Lagergren ER. J Infect Control (Asia) 1:11-30.
The Experimental Animal in Biomedical Research, vol II: Care, Husbandry, and Well-being: An Overview by Species. 1995. Rollin BE, Kesel ML, eds. Boca Raton FL: CRC Press.
UFAW Handbook on the Care and Management of Laboratory Animals, 7th ed, vol 1: Terres trial Vertebrates. 1999. Universities Federation for Animal Welfare. Oxford: Blackwell.

Environmental Enrichment

A novel approach for documentation and evaluation of activity patterns in owl monkeys during development of environmental enrichment programs. 2003. Kondo SY, Yudko EB, Magee LK. Contemp Top Lab Anim Sci 42:17-21.
A review of environmental enrichment for pigs housed in intensive housing systems. 2009. van de Weerd HA, Day JEL. Appl Anim Behav Sci 116:1-20.
A review of environmental enrichment strategies for single-caged nonhuman primates. 1989. Fajzi K, Reinhardt V, Smith MD. Lab Anim 18:23-35.
A targeted approach to developing environmental enrichment for two strains of laboratory F 9778 SpanDC -2.29 -8125 Td [(40.167 0(t)-)]TTjF 9778 >>BDC -2.29 -8225 Td [(A37.917)-151(t)--

Effects of environmental enrichment for mice: Variation in experimental results. 2002. Van de Weerd HA, Aarsen EL, Mulder A, Kruitwagen CL, Hendriksen CF, Baumans V. J Appl Anim Welf Sci 5:87-109.

Effects of environmental enrichment on males of a docile inbred strain of mice. 2004. Marashi V, Barnekow A, Sachser N. Physiol-Behav 82:765-776.

Effects of puzzle feeders on pathological behavior in individually housed rhesus monkeys. 1998. Novak MA, Kinsey JH, Jorgensen MJ, Hazen TJ. Am J Primatol 46:213-227.

Enriching the environment of the laboratory cat. 1995. AWIC Resource Series, No. 2: Envi

Housing, Care and Psychological Well-Being of Captive and Laboratory Primates. 1989. Segal EF, ed. Park Ridge NJ: Noyes Publications.

Incorporation of an enrichment program into a study protocol involving long-term restraint in macaques. 2002. McGuffey LH, McCully CL, Bernacky BJ, Blaney SM. Lab Anim 31:37-39.

Monkey behavior and laboratory issues. 1998. Bayne KA, Novak M, eds. Lab Anim Sci 41:306-359.

Nonhuman Primate Management Plan. 1991. Of ce of Animal Care and Use, NIH. Available at http://oacu.od.nih.gov/regs/primate/primex.htm; accessed January 24, 2010.

Psychological Well-Being of Nonhuman Primates. 1998. National Research Council. Wash ington: National Piona:e(:)-3( )-30(N)--3(s)-3(e)tm; a-4(b)-491(A)-4(n)-4(i)-4(m)-509(S)-4(c)-4(ia(a-4(b-

Joint Working Group on Re nement: Re nement and reduction in production of genetically modi ed mice. 2003. Sixth report of the BVAAWF/FRAME/RSPCA/UFAW Joint Working Group on Re nement. Lab Anim 37(Suppl 1):S1-S49. Available at http://la.rsmjournals.com/cgi/reprint/37/suppl_1/1.pdf; accessed August 10, 2010.

Laboratory animal genetics and genetic quality control. 2003. Festing MF. In: Hau J, Van Hoosier GL Jr, eds. Handbook of Laboratory Animal Science, 2nd ed, vol 1. Boca Raton FL: CRC Press. p 173-203.

Making better transgenic models: Conditional, temporal, and spatial approaches. 2005. Ristevski S. Mol Biotechnol 29:153-163.

Mouse functional genomics requires standardization of mouse handling and housing conditions. 2004. Champy MF, Selloum M, Piard L, Zeitler V, Caradec C, Chambon P, Auwerx J. Mamm Genome 15:768-783.

Persistent transmission of mouse hepatitis virus by transgenic mice. 2001. Rehg JE, Blackman MA, Toth LA. Comp Med 51:369-374.

Projects Involving Genetically Modi ed Animals. 2004. Lane N, Jennings M. RSPCA, Research Animals Department. Available at www.rspca.org.uk/ImageLocator/LocateAsset?asset=documentandassetId=1232712279548andmode=prd; accessed January 24, 2010.

Reproduction and Breeding Techniques for Laboratory Animals. 1970. Hafez E3(p)-3(e)9-3(o)-3(n)-3( )6-3

Domestic Fowl: Good Practice for Housing and Care, 2nd ed. 2008. RSPCA, Research Animals Department. Available at http://content.www.rspca.org.uk/cmsprd/Satellite?blobcol=urldata&blobheader=application%2Fpdf&blobkey=id&blobnocache=false&blobtable=MungoBlobs&blobwhere=1232988515260&ssbinary=true; accessed January 24, 2010.

Ducks and Geese: Good Practice for Housing and Care, 2nd ed. 2008. RSPCA, Research-Animals Department. Available at http://content.www.rspca.org.uk/cmsprd/Satellite?blobcol=urldata&blobheader=application%2Fpdf&blobkey=id&blobnocache=false&blobtable=MungoBlobs&blobwhere=1232988515191&ssbinary=true; accessed January 24, 2010.

Guidelines to the Use of Wild Birds in Research. 2010. Fair J, Paul E, Jones J, eds. Washington: Ornithological Council. Available at http://www.nmnh.si.edu/BIRDNET/guide/index.html; accessed August 20, 2010.

Laboratory Animal Management: Wild Birds. 1977. National Research Council. Washington: National Academy of Sciences.

Laboratory Birds: Re nements in Husbandry and Procedures. 2003. Fifth report of the BVAAWF/FRAME/RSPCA/UFAW Joint Working Group on Re nement. Lab Anim 35(Suppl 1):S1-S163. Available at www.rspca.org.uk/ImageLocator/LocateAsset?asset=document&assetId=1232712322818&mode=prd; accessed August 10, 2010.

Manual of Ornithology: Avian Structure and Function. 2007. Proctor NS, Lynch PJ. In: Thomas NJ, Hunter DB, Atkinson CT, eds. Infectious Diseases of Wild Birds. Oxford: Wiley-Blackwell.

Physiology and Behavior of the Pigeon. 1983. Abs M, ed. London: Academic Press.

Pigeons: Good Practice for Housing and Care, 2nd ed. 2008. RSPCA, Research Animals Department. Available at http://content.www.rspca.org.uk/cmsprd/Satellite?blobcol=

Comfortable quarters for cats in research institutions. 2002. Rochlitz I. In: Comfortable Quarters for Laboratory Animals, 9th ed. Washington: Animal Welfare Institute. p 50-55.

Comfortable quarters for dogs in research institutions. 2002. Hubrecht R. In: Comfortable Quarters for Laboratory Animals, 9th ed. Washington: Animal Welfare Institute. p 56-64.

Dogs: Good Practice for Housing and Care, 2nd ed. 2008. RSPCA, Research Animals Department. Available at http://content.www.rspca.org.uk/cmsprd/Satellite?blobcol=urldata&blobheader=application%2Fpdf&blobkey=id&blobnocache=false&blobtable=MungoBlobs&blobwhere=1232988745038&ssbinary=true; accessed January 24, 2010.

Feline behavioral guidelines from the American Association of Feline Practitioners. 2005. Overall KL, Rodan I, Beaver BV, Careny H, Crowell-Davis S, Hird N, Kudrak S, Wexler-Mitchell E, Zicker S. JAVMA 227:70-84.

Comfortable quarters for nonhuman primates in research institutions. 2002. Reinhardt V. In: Comfortable Quarters for Laboratory Animals, 9th ed. Washington: Animal Welfare

Training nonhuman primates using positive reinforcement techniques. 2003. Prescott MJ, Buchanan-Smith HM, eds. J Appl Anim Welf Sci 6:157-161.
What factors should determine cage sizes for primates in the laboratory? 2004. Buchanan-Smith HM, Prescott MJ, Cross NJ. Anim Welf 13(Suppl):S197-S201.

Rodents and Rabbits

A Laboratory Guide to the Anatomy of the Rabbit, 2nd ed. 1966. Craigie EH. Toronto: University of Toronto Press.
Anatomy and Embryology of the Laboratory Rat. 1986. Hebel R, Stromberg MW. Wörthsee: BioMed Verlag.
Anatomy of the Guinea Pig. 1975. Cooper G, Schiller AL. Cambridge MA: Harvard University Press.
Anatomy of the Rat. 1970. Greene EC. New York: Hafner.
Bensley's Practical Anatomy of the Rabbit, 8th ed. 1948. Craigie EH, ed. Philadelphia: Blakiston.
Biology of the House Mouse. 1981. Symposia of the Zoological Society of London, No. 47. Berry RJ, ed. London: Academic Press.
Chronic stress coping in isolated and socially housed male and female rats. 2003. Westenbroek C, Den B, Gerrits M, Ter H. Society for Behavioral Neuroendocrinology Annual Meeting, Cincinnati, OH.
Comfortable quarters for gerbils in research institutions. 2002. Waiblinger E. In: Comfortable Quarters for Laboratory Animals, 9th ed. Washington: Animal Welfare Institute. p 18-25.
Comfortable quarters for guinea pigs in research institutions. 2002. Reinhardt V. In: Comfortable Quarters for Laboratory Animals, 9th ed. Washington: Animal Welfare Institute. p 38-42.
Comfortable quarters for hamsters in research institutions. 2002. Kuhnen G. In: Comfortable Quarters for Laboratory Animals, 9th ed. Washington: Animal Welfare Institute. p 33-37.
Comfortable quarters for mice in research institutions. 2002. Sherwin CM. In: Comfortable Quarters for Laboratory Animals, 9th ed. Washington: Animal Welfare Institute. p 6-17.
Comfortable quarters for rabbits in research institutions. 2002. Boers K, Gray G, Love J, Mahmutovic Z, McCormick S, Turcotte N, Zhang Y. In: Comfortable Quarters for Laboratory Animals, 9th ed. Washington: Animal Welfare Institute. p 43-49.
Comfortable quarters for rats in research institutions. 2002. Lawlor M. In: Comfortable Quarters for Laboratory Animals, 9th ed. Washington: Animal Welfare Institute. p 26-32.
De nition, Nomenc(m)-3(a)-39a3(t)-3(u)-3(t)-3(e)-3( )-3( )-25(N)Co Td [(D)-3(e)-3((:)-3( )-1o-3( )965 C -3-3(

From house mouse to mouse house: The behavioural biology of free-living Mus musculus and its implications in the laboratory. 2004. Latham M, Mason G. Appl Anim Behav Sci 86:261-289.

Group housing and enrichment cages for breeding, fattening and laboratory rabbits. 1992. Stauffacher M. Anim Welf 1:105.

Group housing f (e)-3(e)-]TJ 2.25 -1.i

Preferences for nesting material as environmental enrichment for laboratory mice. 1997. VandeWeerd HA, van Loo PL, van Zutphen LF, Koolhaas JM, Baumans V. Lab Anim 31:133-143.

Preferences of laboratory mice for characteristics of soiling sites. 1996. Sherwin CM. Anim Welf 5:283-288.

Proceedings of the Third International Workshop on Nude Mice. 1982. Reed ND, ed. vol 1: Invited Lectures/Infection/Immunology; vol 2: Oncology. New York: Gustav Fischer.

Rabbits: Good Practice for Housing and Care, 2nd ed. 2008. RSPCA, Research Animals Department. Available at http://content.www.rspca.org.uk/cmsprd/Satellite?blobcol=urldata&blobheader=application%2Fpdf&blobkey=id&blobnocache=false&blobtable=MungoBlobs&blobwhere=1232988745180&ssbinary=true; accessed January 24, 2010urldata&blobhe

The impact of cage ventilation on rats housed in IVC systems. 2003. Krohn TC, Hansen AK, Dragsted N. Lab Anim 37:85-93.
The impact of low levels of carbon dioxide on rats. 2003. Krohn TC, Hansen AK, Dragsted N. Lab Anim 37:94-99.
The importance of learning young: The use of nesting material in laboratory rats. 2004. van Loo PLP, Baumans V. Lab Anim 38:17-24.
The Laboratory Mouse. 2001. Danneman P, Brayton C, Suckow MA, eds. Boca Raton FL: CRC Press.
The Laboratory Mouse: Selection and Management. 1970. Simmons ML, Brick JO. Englewood Cliffs NJ: Prentice-Hall.
The laboratory rabbit. 1999. Batchelor GR. In: Poole TB, ed. UFAW Handbook on the Care and Management of Laboratory Animals, 7th ed, vol 2. Oxford: Wiley-Blackwell.
The Laboratory Rat, 2nd ed. 2005. Suckow MA, Weisbroth SH, Franklin CL. New York: Aea demic Press. Available at http://198.81.200.2/science/book/9780120749034; accessed January 24, 2010.
The Mouse in Biomedical Research, 2nd ed, vol I: History, Wild Mice, and Genetics. 2007. Fox JG, Barthold SW, Davisson MT, Newcomer CE, Quimby FW, Smith AL, eds. New York: Academic Press.
The Mouse in Biomedical Research, 2nd ed, vol III: Normative Biology, Husbandry, and Models. 2007. Fox JG, Barthold SW, Davisson MT, Newcomer CE, Quimby FW, Smith AL, eds. New York: Academic Press.
The Mouse in Biomedical Research, 2nd ed, vol IV: Immunology. 2007. Fox JG, Barthold SW, Davisson MT, Newcomer CE, Quimby FW, Smith AL, eds. New York: Academic Press.
The Nude Mouse in Experimental and Clinical Research, vol 1, 1978; vol 2, 1982. Fogh J, Giovanella BC, eds. New York: Academic Press.
The Rabbit: A Model for the Principles of Mammalian Physiology and Surgery. 1979. Kaplan HN, Timmons EH. New York: Academic Press.

Other Animals

Aquatic animals as models in biomedical research. 1983. Stoskopf MK. ILAR News 26:22-27.
Captive Invertebrates: A Guide to Their Biology and Husbandry. 1992. Fye FL. Malabar FL: Krieger Publishing.
Effects of experience and cage enrichment on predatory skills of black-footed ferrets (Mustela nigripes). 1999. Vargas A, Anderson SH. J Mammal 80:263-269.
Growth, feed ef ciency and condition of common octopus (Octopus vulgaris) fed on two formulated moist diets. 2008. Cerezo Valverde J, Hernández MD, Aguado-Giménez F, García García B. Aquaculture 275:266-273.
Handbook of Marine Mammals. 1991. Ridgway SH, Harrison RJ, eds. New York: Academic Press.
In uence of diet on growing and nutrient utilization in the common octopus (Octopus vulgaris). 2002. García García B, Aguado-Giménez F. Aquaculture 211:171-182.

Anesthesia and analgesia. 1997. Schaeffer D. In: Kohn DF, ed. Nontraditional Laboratory Animal Species in Anesthesia and Analgesia in Laboratory Animals. San Diego: Academic Press.

Anesthesia and Analgesia in Laboratory Animals. 1997. Kohn DF, Wixson SK, White WJ, Benson GJ, eds. San Diego: Academic Press.

Anesthesia and Analgesia in Laboratory Animals, 2nd ed. 2008. Fish R, Danneman PJ, Brown M, Karas A, eds. 2008. San Diego: Academic Press.

Anesthesia for Veterinary Technicians. 2010. Bryant S, ed. Somerset NJ: Wiley-Blackwell.

Anaesthetic and Sedative Techniques for Aquatic Animals, 3rd ed. 2008. Ross L, Ross B. Somerset NJ: Wiley-Blackwell.

Animal Physiologic Surgery, 2nd ed. 1982. Lang CM, ed. New York: Springer-Verlag.

Textbook of Small Animal Surgery, 3rd ed. 2003. Slatter D. Philadelphia: WB Saunders.
The Biology of Animal Stress: Basic Principles and Implications for Animal Welfare. 2000. Moberg GP, Mench JA. Wallingford UK: CAB International.
The IACUC Handbook. 2006. Flecknell P, Silverman J, Suckow MA, Murthy S, eds. New York: CRC Press.
The importance of awareness for understanding fetal pain. 2005. Lee SJ, Ralston HJ, Drey EA, Partridge JC, Rosen MA. Brain Res Rev 49:455-471.
When does stress become distress? 1999. Moberg GP. Lab Anim 28:422-426.

Disease Surveillance, Diagnosis, and Treatment

Clinical Laboratory Animal Medicine. 2008. Hrapkiewicz K, Medina L, eds. San Francisco: Wiley-Blackwell.
Current strategies for controlling/eliminating opportunistic microorganisms. 1998. White WJ, Anderson LC, Geistfeld J, Martin D. ILAR J 39:391-305.
Drug Dosage in Laboratory Animals: A Handbook. 1989. Borchard RE, Barnes CD, Eltherington LG. West Caldwell NJ: Telford Press.
FELASA recommendations for the health monitoring of breeding colonies and experimental units of cats, dogs and pigs. 1998. Rehbinder C, Baneux P, Forbes D, van Herck H, Nicklas W, Rugaya Z, Winkler G. Report of the Federation of European Laboratory Animal Science Associations (FELASA) Working Group on Animal Health. Lab Anim 32:1-17.
Ferrets, Rabbits and Rodents: Clinical Medicine and Surgery. 1997. Hillyer EV, Quesenberry KE. Philadelphia: WB Saunders.
Handbook of Veterinary Drugs: A Compendium for Research and Clinical Use. 1975. Rossoff IS. New York: Springer.
Invertebrate Medicine. 2006. Lewbart GA. Ames: Blackwell Publishing.
Kirk and Bistner's Handbook of Veterinary Procedures and Emergency Treatment, 8th ed. 2006. Ford RB, Mazzaferro E. Philadelphia: WB Saunders.
Laboratory Animal Medicine. 2002. Fox JG, Anderson LC, Loew FM, Quimby FW, eds. New York: Academic Press.
Mosby's Fundamentals of Animal Health Technology: Principles of Pharmacology. 1983. Giovanni R, Warren RG, eds. St. Louis: CV Mosby.
Veterinary Applied Pharmacology and Therapeutics, 5th ed. 1991. Bradley GC, Pugh DM, Bywater RJ. London: Baillere Tindall.

Blood: Atlas and Sourcebook of Hematology, 2nd ed. 1991. Kapff CT, Jandl JH. 1991. Boston: Little and Brown.
Clinical Biochemistry of Domestic Animals, 4th ed. 1989. Kaneko JJ, ed. New York: Academic Press.
Clinical Chemistry of Laboratory Animals. 1988. Loeb WF, Quimby FW. New York: Pergamon Press.
Color Atlas of Comparative Veterinary Hematology. 1989. Hawkey CM, Dennett TB. Ames: Iowa State University Press.
Color Atlas of Hematological Cytology, 3rd ed. 1992. Hayhoe GFJ, Flemans RJ. St. Louis: Mosby Year Book.
Comparative Neuropathology. 1962. Innes JRM, Saunders LZ, eds. New York: Academic Press.
Duncan and Prasse's Veterinary Laboratory Clinical Pathology. 2003. Latimer KS, Mahaffey EA, Prasse KW. San Francisco: Wiley Blackwell.
Essentials of Veterinary Hematology. 1993. Jain NC. Philadelphia: Lea and Febiger.
Flynn's Parasites of Laboratory Animals, 2nd ed. 2007. Baker DG, ed. Ames: Iowa State University Press.
Handbook of Laboratory Animal Bacteriology. 2000. Hanson AK. Boca Raton FL: CRC Press.
Immunologic Defects in Laboratory Animals. 1981. Gershwin ME, Merchant B, eds. New York: Plenum.
Laboratory Pro les of Small Animal Diseases. 1981. Sodikoff C. Santa Barbara: American Veterinary Publications.
Natural Pathogens of Laboratory Animals: Their Effects on Research. 2003. Baker DG. Washington: American Society for Microbiology.
Pathology of Domestic Animals, 4th ed. 1992. Jubb KVF, Kennedy PC, Palmer N, eds. 1992. New York: Academic Press.
Pathology of Laboratory Animals. 1978. Benirschke K, Garner FM, Jones TC. 1978. New York: Springer-Verlag.
The Pathology of Laboratory Animals. 1965. Ribelin WE, McCoy JR, eds. Spring eld IL: Charles C Thomas.
Veterinary Clinical Parasitology, 6th ed. 1994. Sloss MW, Kemp RL. 1994. Ames: Iowa State University Press.
Veterinary Pathology, 5th ed. 1983. Jones TC, Hunt RD. Philadelphia: Lea and Febiger.

Species-Speci c References—Veterinary Care

Agricultural Animals

Basic Surgical Exercises Using Swine. 1983. Swindle MM. 1983. New York: Praeger.
Diseases of Poultry, 9th ed. 1991. Calnek BW, Barnes HJ, Beard CW, Reid WM, Yoder HW, eds. Ames: Iowa State University Press.
Diseases of Sheep. 1974. Jensen R. Philadelphia: Lea and Febiger.
Diseases of Swine, 7th ed. 1992. Leman AD, Straw BE, Mengeline WL, eds. Ames: Iowa State University Press.
FELASA recommendations for the health monitoring of breeding colonies and experimental units of cats, dogs and pigs. 1998. Rehbinder C, Baneux P, Forbes D, van Herck H, Nicklas W, Rugaya Z, Winkler G. Report of the Federation of European Laboratory Animal Science Associations (FELASA) Working Group on Animal Health. Lab Anim 32:1-17.
Swine in the Laboratory: Surgery, Anesthesia, Imaging, and Experimental Techniques, 2nd ed. Swindle MM, ed. 2007. Boca Raton FL: CRC Press.

Techniques in Large Animal Surgery, 3rd ed. 2007. Hendrickson D, ed. 2007. Somerset NJ: Wiley-Blackwell.
Textbook of Large Animal Surgery, 2nd ed. 1987. Oehme FW, Prier JE. Baltimore: Williams and Wilkins.

The ability of clove oil and MS-222 to minimize handling stressing rainbow trout (Oncorhynchus mykiss Walbaum). 2003. Wagner GN, Singer TD, McKinley RS. Aquacult Res 34:1139-1146.
The ef cacy of clove oil as an anaesthetic for rainbow trout,Oncorhynchus mykiss(Walbaum). 1998. Keene JL, Noakes DLG, Moccia RD, Soto CG. Aquacult Res 29:89-101.
The neurobehavioral nature of shes and the question of awareness and pain. Rose JD. 2002. Rev Fish Sci 10:1-38.
The Pathology of Fishes. 1975. Ribelin WE, Migaki G, eds. Madison: University of Wisconsin.

Birds

Zoo and Wild Animal Medicine Current Therapy, 6th ed. 2007. Fowler ME, Miller RE, eds. Philadelphia: WB Saunders.
Zoo Animal and Wildlife Immobilization and Anesthesia. 2007. West G, Heard D, Caulkett N, eds. Oxford: Wiley-Blackwell.

Nonhuman Primates

Handbook of Primate Husbandry. 2005. Wolfensohn S, Honess P. Oxford: Blackwell
Nonhuman Primates in Biomedical Research: Diseases. 1988. Bennett BT, Abee CR, Henrickson R, eds. New York: Academic Press.
Nursery Rearing of Nonhuman Primates in the 21st Century. 2005. Sackett GP, Ruppenthal GC, Elias K. Chicago: University of Chicago Press.
Primate Parasite Ecology: The Dynamics and Study of Host-Parasite Relationships. 2009. Huffman MA, Chapman CA, eds. West Nyack NY: Cambridge University Press.
Simian Virology. 2009. Voevodin AF, Marx PA. Ames IA: Wiley-Blackwell.
The Emergence of Zoonotic Diseases. 2002. Burroughs T, Knobler S, Lederberg J, eds. Institute of Medicine. Washington: National Academies Press.
Zoonoses and Communicable Diseases Common to Man and Animals. 2003. Acha PN, Szyfres B. Washington: Pan American Health Organization.

Rodents and Rabbits

A Guide to Infectious Diseases of Guinea Pigs, Gerbils, Hamsters, and Rabbits. 1974. National Research Council. Washington: National Academy of Sciences.
Anesthesia and analgesia for laboratory rodents. 2008. Gaertner D, Hallman T, Hankenson F, Batchelder M. In: Fish R, Brown M, Danneman P, Karas A, eds. Anesthesia and Analgesia in Laboratory Animals. San Diego: Academic Press. p 239-298.
Aversion to gaseous euthanasia agents in rats and mice. 2002. Leach MC, Bowell VA, Allan TF, Morton DB. Comp Med 52:249-257.
Behavioural and cardiovascular responses of rats to euthanasia using carbon dioxide gas. 1997.

Eradication of infection with Helicobacter spp. by use of neonatal transfer. 2000. Truett GE, Walker JA, Baker DG. Comp Med 50:444-451.

Eradication of murine norovirus from a mouse barrier facility. 2008. Kastenmayer RJ, Perdue KA, Elkins WR. JAALAS 47:26-30.

Euthanasia of neonatal mice with carbon dioxide. 2005. Pritchett K, Corrow D, Stockwell J, Smith A. Comp Med 55:275-281.

Euthanasia of rats with carbon dioxide: Animal welfare aspects. 2000. Hackbarth H, Kuppers N, Bohnet W. Lab Anim 34:91-96.

Pathology of Aging Syrian Hamsters. 1983. Schmidt RE, Eason RL, Hubbard GB, Young JT, Eisenbrandt DL. Boca Raton FL: CRC Press.

Pathology of Laboratory Rodents and Rabbits, 3rd ed. 2008. Percy DH, Barthold SW. San Francisco: Wiley-Blackwell.

Pathology of the Syrian Hamster. 1972. Homburger F, Baker NY, Kagan E. Basel NY Karger.

Preliminary recommendations for health monitoring of mouse, rat, hamster, guinea pig, gerbil and rabbit experimental units. 1995. Hem A, Hansen AK, Rehbinder C, Voipio HM, Engh E. Scand J Lab Anim Sci 22:49-51.

Prenatal transmission and pathogenicity of endogenous ecotropic murine leukemia virus AKV. 1999. Hesse I, Luz A, Kohleisen B, Er e V, Schmidt J. Lab Anim Sci 49:488-495.

Design and optimization of air ow patterns. 1994. Reynolds SD, Hughes H. Lab Anim 23:46-49.

Design of Biomedical Research Facilities. 1981. Proceedings of the National Cancer Institute Symposium, National Cancer Institute. Monograph Series, vol 4. NIH Pub. No. 81-2305.

Design of surgical suites and post surgical care units. 1997. White WJ, Blum JR. In: Kohn DF, Wixson SK, White WJ, Benson GJ, eds. Anesthesia and Analgesia in Laboratory Animals. San Diego: Academic Press.

Estimating heat produced by laboratory animals. 1964. Brewer NR. Heat Piping Air Cond 36:139-141.

Guidelines for Construction and Equipment of Hospitals and Medical Facilities, 2nd ed. 1987. American Institute of Architects Committee on Architecture for Health. Washington: American Institute of Architects Press.

Guidelines for Laboratory Design: Health and Safety Considerations. 1993. DiBerardinis LJ, Baum JS, First MW, Gatwood GT, Groden EF, Seth AK. New York: John Wiley and Sons.

Handbook of Facilities Planning, vol 2: Laboratory Animal Facilities. Ruys T, ed. New York: Van Nostrand.

Laboratory Animal Houses: A Guide to the Design and Planning of Animal Facilities. 1976. Clough G, Gamble MR. LAC Manual Series No. 4. Carshalton UK: Laboratory Animals Centre.

Laboratory Animal Housing. 1978. National Research Council. Washington: National Academy of Sciences.

Livestock behavior and the design of livestock handling facilities. 1991. Grandin T. In: Ruys T, ed. Handbook of Facilities Planning, vol 2: Laboratory Animal Facilities. New York: Van Nostrand. p 96-125.

Management and Design: Breeding Facilities. 2007. White WJ. In: Fox JG, Barthold SW, Davison MT, Newcomer CE, Quimby FW, Smith AL, eds. The Mouse in Biomedical Research, 2nd ed, vol III: Normative Biology, Husbandry, and Models. New York: Academic Press. p 235-269.

Planning and Designing Animal Research Facilities. 2008. Hessler J, Lehner N, eds. Orlando: Academic Press.

Rodent Facilities and Caging Systems. 2009. Lipman NS. In: Hessler J, Lehner N, eds. Planning and Designing Animal Research Facilities. Orlando: Academic Press. p 265-288.

Structures and Environment Handbook, 11th ed, rev 1987. Ames: Midwest Plan Service, Iowa State University.

Warning! Nearby construction can profoundly affect your experiments. 1999. Dallman MF, Akana SF, Bell ME, Bhatnagar S, Choi SJ, Chu A, Gomez F, Laugero K, Sorian L, Viau V. Endocrine 11:111-113.

Working safely at animal biosafety level 3 and 4: Facility design and management implications. 1997. Richmond JY, Ruble DL, Brown B, Jaax GP. Lab Anim 26:28-35.

APPENDIX

B

U.S. Government Principles for the Utilization and Care of Vertebrate Animals Used in Testing, Research, and Training

The development of knowledge necessary for the improvement of the health and well-being of humans as well as other animals requires in vivo experimentation with a wide variety of animal species. Whenever U.S. Government agencies develop requirements for testing, research, or training procedures involving the use of vertebrate animals, the following principles shall be considered; and whenever these agencies actually perform or sponsor such procedures, the responsible Institutional Official shall ensure that these principles are adhered to:

I. The transportation, care, and use of animals should be in accordance with the Animal Welfare Act (7 U.S.C. 2131 et. seq.) and other applicable Federal laws, guidelines, and policies.[1]

II. Procedures involving animals should be designed and performed with due consideration of their relevance to human or animal health, the advancement of knowledge, or the good of society.

III. The animals selected for a procedure should be of an appropriate species and quality and the minimum number required to obtain valid results. Methods such as mathematical models, computer simulation, and in vitro biological systems should be considered.

[1]For guidance throughout these Principles, the reader is referred to the Guide for the Care and Use of Laboratory Animals prepared by the Institute for Laboratory Animal Research, National Academy of Sciences.

IV. Proper use of animals, including the avoidance or minimization of discomfort, distress, and pain when consistent with sound scienti c practices, is imperative. Unless the contrary is established, investiga tors should consider that procedures that cause pain or distress in human beings may cause pain or distress in other animals.

V. Procedures with animals that may cause more than momentary or slight pain or distress should be performed with appropriate sedation, analgesia, or anesthesia. Surgical or other painful procedures should not be performed on unanesthetized animals paralyzed by chemical agents.

VI. Animals that would otherwise suffer severe or chronic pain or distress that cannot be relieved should be painlessly killed at the end of the procedure or, if appropriate, during the procedure.

VII. The living conditions of animals should be appropriate for their spe cies and contribute to their health and comfort. Normally, the hous ing, feeding, and care of all animals used for biomedical purposes

APPENDIX
C

Statement of Task

The use of laboratory animals for biomedical research, testing and education is guided by the principles of the Three Rs, replacement of animals where acceptable non-animal models exist, reduction in the number of animals to the fewest needed to obtain statistically s(y)-3nisti3(c)-3(t

evidence, while maintaining the performance standards of the 1996 *Guide*. The committee will ensure that any recommendations in the *Guide* will be consistent with current Public Health Service Policy, the Animal Welfare Regulations, and the most recent Report of the American Veterinary Medical Association Panel on Euthanasia.

In addition to the published report, the updated *Guide* will be posted on the Internet in a pdf or equivalent format such that users will be able to search the entire document at one time.

APPENDIX
D

About the Authors

Janet C. Garber(Chair), DVM, PhD, received her Doctor of Veterinary Medicine degree from Iowa State University and her PhD in pathophysiology from the University of Wisconsin. Her experiences have included infectious disease research at the U.S. Army Medical Research Institute of Infectious Diseases (USAMRIID), primate medicine and research, GLP device and materials evaluation, and transplantation immunology. Her current interests are in the areas of laboratory animal f3(w)-3(a)-T-3( )-98(i3(m)-3(a)e-3(a)-I32-3
disct 63( )-20( )-20( )-2s raise Il raise resear, (n)-3(o)-3(a)-3(b)-3marast ter
misntiO7Dmt

been associated with IACUCs at small, medium, and large institutions for over two decades and is familiar with the oversight of animal care and use programs. He has served on multiple study sections for both the NIH and DOD. Dr. Barbee also served as an Oxford, UK 2006 fellow (recipient, VCU Harris-Manchester Award) where he examined policies, training, and security issues related to animal care and use within the UK.

Joseph T. Bielitzki, MS, DVM, is Research Manager, University of Central Florida. Dr. Bielitzki has worked with non-human primates in the laboratory environment for 20 years. Over this period he has worked with macaques (pig-tail, long-tail, Japanese, rhesus, stump-tail), baboons (yellow, green and hybrids) squirrel monkeys, capuchin monkeys, mangabeys, gibbons, chimpanzees, orangutans, bonobos, and gorillas. In the area of non-human primates his area of expertise is in enteric diseases, nursery rearing, and colony management. He has also worked with mice and rats in a variety of international facilities. He was instrumental in the writing

of Laboratory Animal Science. He began his career in the pharmaceutical industry in 1994 as Senior Director of Worldwide Laboratory Animal Resources for Rhone Poulenc-Rorer Pharmaceuticals, leading to his position as Vice President of Laboratory Animal Science and Welfare for Aventis Pharmaceuticals in 1999. In 2001 he moved to Wyeth Pharmaceuticals where he was Vice President of BioResources until his retirement in 2007. During his career, Dr. Donovan served in numerous professional leadership positions including President of ACLAM and President of the Board of Directors of the Pennsylvania Society for Biomedical Research (PSBR). He also served on several boards including those of the National Association for Biomedical Research, ACLAM, PSBR, and the New Jersey Association for Biomedical Research.

Dennis F. Kohn, DVM, PhD, is Professor Emeritus of Clinical Comparative Pathology at Columbia University. He received his DVM from Ohio State University and a doctorate in Medical Microbiology from West Virginia University. He is board certified by the American College of Laboratory Animal Medicine. He has directed laboratory animal resource/comparative medicine programs at West Virginia University Medical Center, University of Texas Medical School at Houston, and the Health Sciences Division, Columbia University. His research interests have dealt primarily with the pathogenicity of Mycoplasma pulmonis in the respiratory tract of laboratory rats, and the experimental pathology induced within the CNS and joints of rats inoculated with M. pulmonis. He is a past president of the American College of Laboratory Animal Medicine and the American Society of Laboratory Animal Practitioners, and as chair of a number of American Association of Laboratory Animal Science committees. He has served as a Council member of AAALAC-International, as a member of the 1986 American Veterinary Medical Association's Panel on Euthanasia, and the 1996 Institute of Laboratory Animal Resources' committee to revise the *Guide*.

Neil S. Lipman, VMD, is Director of the Center of Comparative Medicine and Pathology serving the Memorial Sloan-Kettering Cancer Center (MSKCC) and the Weill Medical College of Cornell University and is Professor of Veterinary Medicine in Pathology and Laboratory Medicine at Weill Cornell as well as a Laboratory Member at the Sloan-Kettering Institute at MSKCC. Dr. Lipman is a Diplomate of the American College of Laboratory Animal Medicine with over 25 years' experience in laboratory animal medicine and science. Dr. Lipman has expertise in vivarium design, engineering, and operations, having designed over 1.5 million gross square feet of vivarium space and overseen the operation of a number of major academic animal resource programs. His research interests are principally translational and include development and analysis of new technologies especially with

respect to rodent housing systems and monoclonal antibody production, the characterization of various animal models, understanding the etiopathogenesis of endocrinologic disorders affecting laboratory animal species, and development and analysis of novel therapeutic strategies. Throughout his career, Dr. Lipman has been extensively involved in the postgraduate training of laboratory animal specialists.

Paul Locke,MPH, JD, DrPH, an environmental health scientist and attorney, is Associate Professor at the Johns Hopkins University Bloomberg School of Public Health in the Department of Environmental Health Sciences, Division of Toxicology. He holds an MPH from Yale University School of Medicine, a DrPH from the Johns Hopkins University Bloomberg School of Public Health and a JD degree from Vanderbilt University School of Law. Prior to joining the Department of Environmental Health, he was the Deputy Director of the Pew Environmental Health Commission and the Director of the Center for Public Health and Law at the Environmental Law Institute. Dr. Locke's research and practice focus on how decision makers use environmental health science and toxicology in regulation and policy making and how environmental health sciences in uence the policy-making process. His areas of study include alternatives to animal testing in biomedical research, with particular emphasis on toxicity testing. He also maintains an active research program in radiation studies and radiation protection policy. Dr. Locke directs the Doctor of Public Health degree program in the Department of Environmental Health Sciences and is co-director of the Johns Hopkins certi cate program in Humane Science and Toxicology.

regulations concerning research animal care and use. Her research interests include innate immunity and infectious disease, toxicologic pathology, and the interactions between rodents and their environment as they relate to disease susceptibility. In 2007 she was the inaugural recipient of the North American Animal Welfare Award, co-sponsored by Procter & Gamble and the Humane Society of the United States.

Geoffrey A. Wood, DVM, PhD, DVSc, is Associate Professor in the Department of Pathobiology at the Ontario Veterinary College, University of Guelph, Canada. Dr. Wood has a doctorate in cancer biology and a doctorate in veterinary pathology. He has been involved in design or pathologic characterization of hundreds of genetically engineered rodents, both as the former Associate Director of Pathology at the Centre for Modeling Human Disease in Toronto, and in his current position. His lab conducts research on cancer genetics and the process of metastasis, with a focus on bone and prostate cancer. Dr. Wood's research collaborations include projects investigating various aspects of a wide range of different cancer types, as well as studies on stem cell biology, immunity, and in ammation.

Hanno Würbel, Dr.sc.nat, is Professor of Animal Welfare and Ethology at the Justus Liebig University in Giessen, Germany. He has studied biology (zoology) at the University of Berne, Switzerland and graduated from the ETH Zürich, Switzerland with a doctorate in natural sciences. He has experience in animal behavior and in the scienti c assessment of animal well-being, and has mostly worked with rodents, but also with rabbits, dogs, poultry, and horses. His research focuses on environment-dependent plasticity of brain and behavior in relation to questions of animal husbandry and animal welfare. In 2005 Dr. Würbel received the Hessian Animal Welfare Research Prize and in 2009 the Felix Wankel Animal Welfare Research Award. He is a member of the Animal Welfare Council of the German government, Central Animal Welfare Of cer of the University of Giessen, and head of the University's Central Animal Facility. He is also a council member of the International Society of Applied Ethology (ISAE), editor of the Journal of Applied Animal Welfare Science, and editorial board member of the journal Applied Animal Behaviour Science.

Index

A

Acclimation and adaptation, 111, 188 (see also Procedural habitation and training)
  caloric management and, 67
  to noise and vibrations, 81-82
  to outdoor housing, 44, 54-55
  to physical restraint, 29
Acquisition of animals (see Procurement)
Activity and exercise, xiv, 43, 52-53, 55, 56, 57, 58, 63-64, 67, 84 (see also Enrichment)
Agricultural animals, 2
  biomedical vs. agricultural research, 32-33
  environment and housing, 29, 33, 43, 44, 137, 176-177
  floor drains, 137, 138
  information resources, 33, 176-177, 191-192
  protocol review, 32-33
  space requirements, 60, 62-63
  surgery, 117, 144, 145
  veterinary care, 145, 191-182
Air pressure, 45, 47, 139, 145-146, 149-150
Air quality and airborne contaminants, 45-47
Air flow (see Ventilation and air flow)
Albino animals, 48-49, 141
Allergens and allergic reactions, 18, 19, 20, 22-23, 45, 155
Alternatives to laboratory animal use, 3, 5, 12, 25, 27, 162-167
American Association for Laboratory Animal Science, 16
American College of Laboratory Animal Medicine, 24
  Guidelines for Adequate Veterinary Care, xvii, 14
Ammonia, 45, 47, 68, 70, 71, 72-73, 78, 79, 80, 81, 85
Amphibians, 41-42, 44-45, 46, 77, 78-79, 80, 81, 82, 83, 84, 85, 108, 119, 156, 178-179, 192 (see also Aquatic species)
Anesthesia and analgesia
  considerations in selection of agents, 75, 121
  equipment/delivery, 34, 122, 145, 147
  euthanasia agent, 124
  field conditions, 32, 117
  monitoring, 119-120, 122, 123
  neuromuscular blocking agents, 122-123
  oversight, 34, 156
  pain management, 12, 121-122
  preemptive analgesia, 121
  principles governing use, 12, 14, 26
  recommended reading, 188-190
  recordkeeping, 34

surgery, 115, 116, 117, 118, 119-120, 122-123, 145, 188-190
training, 17, 106, 115, 122
waste gas management, 21, 147
Animal and Plant Health Inspection Service (APHIS), 30, 148
Animal care and use program
attending veterinarian, 14
collaborations, 15
de ned, 6
disaster planning and emergency preparedness, 35
IACUC, 14-15, 24-33
institutional of cial, 13-14
oversight, 24-34
personnel management, 15-24
program management, 13-24
recommended reading, 167-172
regulations, policies, and principles, 12-13
responsibilities, 13-15
Animal care personnel, (see also Personnel)
training and education, 16
Animal training (see Procedural habituation and training)
Animal Welfare Act, 1, 4, 25, 34, 199
Animal Welfare Information Center, xv, 16
Animal Welfare Regulations, 1, 12
housing guidelines, 56, 59
IACUC, 25
transportation of animals, 107
veterinary care, 30
Antibiotics, 116
Anxiolytics, 122-123
Apes (see Nonhuman primates)
Aquatic species, 2 (see also Amphibians; Fish; Reptiles)
algal growth, 86
behavior and social management, 84
diseases, 112
emergency, weekend, and holiday care, 87
enrichment and social housing, 82-83
environment, 77-82
facilities, 150-151
food and feeding, 84-85
housing, 82-83, 150-151
husbandry, 84-87
identi cation, 87
illumination, 81
life support system, 79-80
macroenvironment, 77-78, 86
management, 84-88
microenvironment, 77-78, 82-83
noise and vibration, 81-82
pest control, 87
population management, 87-88
recommended reading, 178-179, 192-193
recordkeeping, 87-88
sanitation, 85-86
sheltered, outdoor, and naturalistic housing, 83
space, 83
substrates, 85
temperature, humidity, and ventilation, 80-81
water quality, 78-79, 85
Aseptic practices, 17, 34, 52, 70, 75, 115, 116, 117, 118-119, 144
Attending veterinarian, 3, 4, 13, 14, 15, 17, 21, 24, 28, 106, 114, 116 (see also Veterinary care)
Autoclaving
bedding material, 69, 146
drinking water, 146
feed, 66, 146
surgical instruments, 119, 145
Avian species (see Birds)

B

Behavioral and social management
acclimation, 111
activity and exercise, 63-64, 84
air ow and ventilation and, 150
aquatic species, 82, 83, 84, 85
bedding, nesting, and substrate materials and, 49, 52, 85
enrichment, 52-53, 54, 60, 83, 85
food and feeding, 65
illumination and, 47, 48, 49
procedural habituation and training, 29, 64-65
recordkeeping, 75
sanitation and, 72
social environment, 51, 58, 59, 64, 111
space considerations, 55, 56, 57, 58, 59
temperature and humidity and, 43, 44, 80
terrestrial species, 83-85
training of personnel, 53, 121
veterinary care and, 107, 111, 112, 114
Behavioral changes
infections and, 118
monitoring animals for, 112
moribund state, 28
pain and distress, 120-121, 123
physical restraint and, 29
sanitation and, 71
stereotypies and compulsive behaviors, 63
transportation of animals and, 150
Behavioral research, 30, 31, 111, 135, 149-150
Bioexclusion, 28 (see alsoBarriers)
Biologic agents and hazards, 18, 19, 20, 21, 111, 112-113, 148, 170-172
Biosecurity, animal, xviii, 105, 107, 108, 109-110, 111, 135, 170-172
Birds, 41, 44, 48, 49, 60, 121, 144, 156, 179-180, 193
Bites and scratches, 18, 23
Breeding and reproduction
aquatic animals, 44, 80, 82, 83, 87, 88
caloric restriction and, 67
cryopreservation of fertilized embryos, ova, ovaries, or spermatozoa, 76, 107
enriched environments and, 83
genetic management, 76, 77
GMAs, 28, 76
lighting and, 48, 137
nesting, bedding and substrates, 52, 70, 83
nomenclature, 77
nonhuman primates, 59
recordkeeping, 76, 77, 87, 88
rodents and rabbits, xviii, 57, 77, 107
sanitation activities and, 70
space requirements, 55, 56, 57, 33
temperature and humidity and, 44, 80, 82
vibration and, 50
Building Seismic Safety Council, 133

C

Cages and caging(see also Housing; Microenvironment)
cleaning and sanitation, 18, 46, 47, 52, 69, 70-71, 72, 73, 86, 142, 143, 145
dimensions and space allocation, xviii, 56, 57, 58, 59, 60, 61, 133
dividers and shelters in, 53
ooring, 70
isolators, 45, 47, 52, 111, 139, 146
light intensity in, 46, 49
location on cage rack, 46, 49
mechanical washers, 71, 72, 86, 135, 136, 141, 143
metabolic, 121
in secondary enclosures, 47
ventilated, 44, 46, 47, 50, 58, 70-71, 111, 141, 146
vibration of, 50
Caloric management, 31, 57, 67
Canadian Council on Animal Care, 16
Carcass disposal, 20, 73-74, 138, 142
Cardiovascular shock models, 27
Cats
behavioral and social management, 58, 59, 63-64, 180-181
enrichment, 53
environmental requirements, 43, 44, 53, 180-181
housing and space, 49, 56, 58, 59, 180-181
procurement, 106
recommended reading, 180-181, 193
recordkeeping, 75-76
veterinary care, 193
Cattle, 62-63, 64, 176-177, 191-192 (see also Agricultural animals)
Cedar shavings, 69, 156

Ceilings, 56, 81, 142
Centers for Disease Control and Prevention, 19, 21, 107, 148, 157
Cercopithecine herpesvirus 1, 23
Chairing, 29
Chemical agents
  cleaning materials, 71, 72
  non-pharmaceutical-grade, 31
  hazardous, 18, 19, 20, 21, 136
Chickens, 48, 60
Chimpanzees, xviii, 61
Chlorines and chloramines, 72, 78-79, 80, 85, 86
Circadian rhythm, 43, 48, 49, 137, 149
Cleaning agents, 18, 71, 72 (see also Sanitation)
  (see also Veterinary care)
Clinical care and management
  emergency care, 114
  medical management, 114
  recordkeeping, 115
Clothing and footwear, 20, 21, 146
Cold storage and refrigeration, 66, 73, 136, 142
Cold stress, 43, 44, 53
Collaborations, interinstitutional, 15
Computer modeling
  alternative to animal use, 5, 12, 25, 199
  facility design, 46, 133
  genetic variability, 76
Conditioned-response protocols, 31
Construction guidelines, 136-143
Containment of hazardous materials, 21, 148-149
Contaminants
  airborne, 22, 45, 46-47, 139, 140, 143, 145, 146-147, 149
  bedding, 69, 141, 146
  cleaning implements, 72, 85
  food, 19, 65, 66, 67, 84, 141, 146
  genetic, 76
  microbial, 72, 81, 108, 109, 110, 113, 118, 144-145
  recommended readings
  water, 67-68, 81, 86, 138
Convention on International Trade in Endangered Species of Wild Fauna and Flora, 107
Corridors, 86, 134, 136, 137, 141, 142
Cryogens, 147
Cryopreservation, 76, 107
Cyanobacteria, 86

D

Defecation, 72
De nitions
  animal biosecurity, 109
  animal care and use program, 6, 11
  animal use, 2
  attending veterinarian, 14
  engineering standard, 6
  humane care, 6
  institutional animal care and use committee, 14
  institutional of cial, 13
  laboratory animals, 2
  life support system, 79
  macroenvironment, 42
  “may,” 8
  microenvironment, 42
  “must,” 8
  performance standard, 6-7
  policies, 7
  practice standard, 7
  principles, 7
  procedures, 7
  “should,” 8
  thermoneutral zone, 43
  three Rs, 4-5
Dehydration, 68
Desiccation stress, 45, 46
Diet and dietary control (see also Food and feedng)
  abrupt changes in, 67
  aquatic animals, 84, 85
  autoclavable, 66
  caloric management, 31, 57, 67
  certi ed diets, 65
  chemically de ned diets, 65, 66
  natural-ingredient, 65, 66
  nutritionally balanced, 67
  puri ed diets, 65, 66
  quality assurance, 65, 66
  treats, 67
Disaster planning, 35, 74-75
Disease control, 23, 72, 81, 108, 109, 110, 112-113, 118, 144-145, 190
Disease susceptibility, 19, 42

Disinfection, 20, 51, 52, 68, 69, 70-72, 78, 79, 80, 84, 86, 110, 118, 119, 138, 143, 144, 150
Disposable clothing, 146
Distress (see also Pain and distress)
  de ned, 121
  neuromuscular blocking agents and, 122-123
  recommended reading, 188-190
Diurnal cycles, 48, 141
Dogs
  behavioral and social management, 29, 59, 63-64, 180-181
  enrichment, 53
  environmental requirements, 43, 44, 180-181
  housing and space, 49, 58, 59, 136, 138, 180-181
  infusion pumps, 29
  pain and distress, 121
  procurement, 106
  recommended reading, 180-181, 193
  recordkeeping, 75-76
  veterinary care, 193
Doors, 136, 137, 138, 142, 143, 144, 149, 150
Drafts, 46, 140
Drainage, 54, 136, 138, 143, 144, 149, 150

E

Ecdysis (molting), 45, 156

Hydrogen peroxide, 72
Hygiene, personal, 18, 19, 20

I

Identi cation of animals, 75, 87
Illumination/lighting, 47-49, 81, 141
Imaging facilities, 146-147
Immunization, 22
Immunode cient animals, 21, 28, 66, 145
Infectious agents (see Biologic agents and hazards
Infectious disease research, 27
Inspection of facilities and equipment, 34
Institute for Laboratory Animal Resources (ILAR), 2, 16
Institutional Animal Care and Use Committee
  agricultural animals, 32-33
  animal care and use programs and, 6, 11, 13
  collaborating institutions, 15
  endpoint considerations, 27-28
  eld investigations, 32
  food and uid regulation, 30-31
  investigating and reporting animal welfare concerns, 23
  membership and function of committee, 17, 24-25
  multiple survival surgical procedures, 30
  non-pharmaceutical-grade chemicals or substances, 31
  occupational health and safety and, 17
  physical restraint, 29-30
  program management and, 13, 14-15
  protocol review, 25-26
  recommended reading, 169-170
  special considerations during review process, 27-33
  surgery and, 115-118
  surgical procedures, 30
  training and education, 15
  unexpected outcomes, 28-29
Institutional of cial, 13-14, 17, 23, 24, 25, 30, 35
International Air Transport Association, 107
International Committee on Standardized Genetic Nomenclature for Mice, 77
Invertebrate animals, 2
Irradiation facilities, 157-158

L

Laboratory Animal Welfare and Training Exchange, 16
Latex toxicity, 84
Laws, regulations, and policies (see also Animal Welfare Act; Animal Welfare Regulations; Public Health Service Policy; other speci c acts and policies)
  recommended reading, 168
  training in, 17
Lighting (see Illumination/lighting)
Lymphocytic choriomeningitis virus, 113

M

Macacine herpesvirus, 23, 112
Macaques, 23, 112
Macroenvironment
  aquatic species, 77-78, 86
  de ned, 42
  terrestrial species, 42
Management of animals (see Behavioral and social management; Husbandry; Population management)
  aquatic species, 84-88
  medical, 114
  recommended reading, 172-173
  terrestrial species, 63-77
Mice (see also Rodents)
  acclimation, 111
  aggression in males, 53, 64
  anxiety studies, 53
  bedding and nesting materials, 52, 53, 68
  breeding animals, 52
  caloric restriction, 67
  enrichment, 53, 54
  GMAs, 28, 68, 145-146
  housing, 53, 57, 145-146
  identi cation/nomenclature, 77
  illumination, 49, 137
  infections, 111, 112
  nude or hairless animals, 68
  recommended reading, 184-187, 194-196
  social stress, 53
  space requirements, 57, 156
  speci c pathogen-free mice, 145-146
  temperature and humidity, 43, 45
  toe-clipping, 75

Microenvironment (primary enclosure)
aquatic species, 77-78, 82-83
de ned, 42
terrestrial species, 42, 50-52
Midwest Plan Service, 33
Monitoring and surveillance
disease prevention, 112-113
facilities, equipment, and exposure, 19-20
food and uid intake, 31
GMAs, 28
intraoperative, 119
postapproval, 33-34
postoperative care, 119

recommended reading, 169-170
responsibility for, 13
Owl monkeys (*Aotus trivirgatus*), 112

P

Pain and distress (see also Distress)
analgesics and anesthetics, 12, 121-123
chronic, 122
guidelines and, 12, 199-200
humane endpoints, 5, 27, 123
indicators of, 120-121, 123
modeling/studies, 28
monitoring GMAs for, 29
multiple survival surgical procedures, 30
postoperative, 117, 120
prevention or alleviation, 5, 12, 26, 33, 35, 105, 114, 120-121, 200
recommended reading, 120, 188-190
restraint devices, 29
studies of, 34
"wind-up" phenomenon, 120
Parvoviruses, 113
Pasteurella multocida, 112
Pedigree management, 55, 76, 87
Performance standard, de ned, 6-7
Personnel (see also Occupational health and safety program; Training and education; speci c personnel)
animal welfare concerns and, 23-24
hygiene, 18, 19, 20
management, 15-24
medical evaluation and preventive medicine for, 22-23
occupational health and safety program, 17-23
protection, 21-22
security, 23
Pest control, 74, 87, 110
Pesticides, 65, 74
Photoperiod, 47, 48, 81, 137, 143
Photostressors, 47s (see also Illumination/lighting)
Phototoxic retinopathy, 48, 49
Physical hazards, 18, 19
Physical plant (see also Facilities)

Protocols for animal use
alternatives to animal use, 3, 5, 12, 25, 27, 162-164
conditioned-response, 31
food and fluid regulation, 30-31
IACUC review, 26-33
number of animals used, 5, 12, 15, 25, 26, 28, 106, 199, 201
postapproval monitoring, 33-34
reuse of animals, 5, 30
special considerations, 27-33
Three Rs, xvii, 3, 4-5, 17, 201
topics considered, 25-26
Public Health Service Policy on Humane Care and Use of Laboratory Animals, 1, 12, 25, 56

Q

Quail, 60, 180, 193
Quarantine and stabilization, 110-111

R

Rabbits
behavioral and social management, 63
enrichment, 54
environment, 43, 44, 49, 54, 140
food and feeding, 67
housing, 51, 53, 58, 59
infections, 112
nomenclature, 77
recommended reading, 184-187
sanitation of cages, 71 n.2
space requirements, xviii, 59
Rabies, 22
Radiation hazards, 18, 20, 73, 147
Rainbow trout, 112
Rat Genome and Nomenclature Committee, 77
Rat theilovirus, 112
Rats (see also Rodents)
acclimation, 111
albino, 48-49
bedding and nesting materials, 52
breeding animals, 52
caloric restriction, 67
diseases, 45, 111, 112
GMAs, 145-146
identification/nomenclature, 77
illumination, 48-49, 137
recommended reading, 184-187, 194-196
space requirements, 57, 156
temperature and humidity, 43, 44, 45
Recordkeeping
clinical care, 115
population management, 75-76, 87-88
Removal of animals from protocol, 26, 29, 31
Reporting
accidents, bites, scratches, and allergic reactions, 23
animal welfare concerns, 17, 23-24
hazardous conditions or “near miss” incidents, 19
status of Animal care and use program, 25
Reproduction (see Breeding and reproduction)
Reptiles, 41-42, 44, 45, 77, 80, 81, 82, 83, 84, 85, 156, 178-179, 192-193
Research and testing methodology, recommended reading, 165-166
Research team, training and education, 16-17
Respirators and respiratory protection, 22, 23
Restraint, 29-30, 55, 122, 144
Reuse of animals, 5
Ringtail, 45
Rodents, laboratory (see also individual species)
acclimation, 111
aggression in males, 53, 64
albino animals, 48-49, 141
anxiety studies, 53
barrier facilities, 145-146
bedding and nesting materials, 52, 53, 68-69
breeding, xviii, 49, 52, 57
caloric restriction, 67
diseases, 111, 112, 113
enrichment, 53, 54
euthanasia, 124
GMAs, 28, 68, 145-146
housing, 53, 57, 145-146
identification/nomenclature, 77
illumination, 48, 49, 137
implanted systems, 29
noise and vibration effects, 49, 50

pain and distress, 121-122
recommended reading, 184-187, 194-196
social stress, 53
space requirements, xviii, 56-58, 156
temperature and humidity, 43, 45
thermoneutral zone, 43

S

Sanitation

Training and education, 20, 115-116
animal care personnel and technicians, 16, 50, 78
aquatic systems, 78
emergency preparedness, 35
IACUC members, 17
IACUC oversight of, 15
noise reduction, 50
postapproval review as, 34
recommended reading, 168-169
research team, 16-17
U.S. Government Principles, 12, 199-200
veterinary and other professional staff, 15-16
Transportation of animals, 12, 14, 32, 105, 107-109, 110, 111, 134, 135, 147, 150, 156, 188, 199
Trauma research, 27
Tuberculosis, 23, 112
Tumor models/studies, 27, 34

U

Unexpected outcomes, 28-29, 34
US Department of Agriculture, 19, 30, 107, 148, 157
US Fish and Wildlife Service, 106, 107, 157
U.S. Government Principles for the Utilization and Care of Vertebrate Animals Used in Testing, Research, and Training, 4, 12, 199-200
Uses and users of the Guide, 4-5

V

Vaccine challenges/studies, 27
Ventilation and air ow (see also Air qual 3(l)-3(a)-3(t)-3(EMC pT5)-3(6)-3(,)-3( )-31(1)-3(8)-3(8)-3(,)-3( )-6es

VTJ EM0-3(96DC -3 -1<</MC0-3(9n)-3(t)-3(i)-3(l)-3(a)-3(t)-357)-3(c
ses a d us Vo-9-9 Ai 9(96 19-